

Product Datasheet

beta-Catenin [p Ser552] Antibody - BSA Free NBP3-05441

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-05441

Updated 2/14/2024 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-05441



NBP3-05441

beta-Catenin [p Ser552] Antibody - BSA Free


Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Polyclonal
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS with 50% glycerol, pH7.3.

Product Description	
Host	Rabbit
Gene ID	1499
Gene Symbol	CTNNB1
Species	Human, Mouse, Rat
Immunogen	A synthetic phosphorylated peptide around S552 of human beta-Catenin (NP_001091679). RTSMG


Product Application Details	
Applications	Western Blot
Recommended Dilutions	Western Blot 1:500-1:2000

Images

Western Blot: beta-Catenin [p Ser552] Antibody [NBP3-05441] - Western blot analysis of extracts of C2C12 cells, using beta-Catenin pAb (NBP3-05441) at 1:1000 dilution or beta-Catenin antibody (A11932). C2C12 cell lysates were treated by CIP at 37C for 1 hour. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Enhanced Kit. Exposure time: 30s.



Western Blot: beta-Catenin [p Ser552] Antibody [NBP3-05441] - Western blot analysis of extracts of various cell lines, using beta-Catenin [p Ser552] antibody (NBP3-05441) at 1:1000 dilution. C2C12 cell lysates were treated by CIP at 37C for 1 hour. NIH/3T3 cells were treated by Forskolin (30uM) for 30 minutes after serum-starvation overnight or treated by CIP at 37C for 1 hour. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% BSA. Detection: ECL Enhanced Kit. Exposure time: 30s.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA

Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada

Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420

info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-05441

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-61628PEP	beta-Catenin Recombinant Protein Antigen

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-05441

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

