## **Product Datasheet**

# Bromodeoxyuridine/BrdU Antibody (BRDU/3902R) NBP3-07261-100ug

Unit Size: 100 ug Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-07261

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-07261



## NBP3-07261-100ug

Decree Level 2 Law (D. H. A. Clark, (D.D.H. (2000D))		
Bromodeoxyuridine/BrdU Antibody (BRDU/3902R)		
Product Information		
Unit Size	100 ug	
Concentration	0.2 mg/ml	
Storage	Store at 4C.	
Clonality	Monoclonal	
Clone	BRDU/3902R	
Preservative	0.05% Sodium Azide	
Isotype	IgG	
Purity	Protein A or G purified	
Buffer	10 mM PBS with 0.05% BSA	
Product Description		
Description	200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08948)  Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C.	
Host	Rabbit	
Species	Non-species specific	
Marker	Proliferation Marker	
Specificity/Sensitivity	It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting Sphase cells, providing useful information on the aggressiveness of tumors.	
Immunogen	Bromodeoxyuridine/BrdU conjugated to KLH	
Product Application Details		
Applications	Immunohistochemistry-Paraffin	
Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml	
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. For	

<b>Product Application Details</b>	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. For staining of formalin-fixed tissues, incubate sections in 4N HCl for 30 minutes at RT followed by digestion with trypsin at 1mg/ml PBS, 10 min at 37C. Optimal dilution for a specific application should be determined.





#### **Novus Biologicals USA**

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP3-07261-100ug

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP3-07125 Bromodeoxyuridine/BrdU MCF-7 Cell Line Slides (Adult

Adenocarcinoma)- Paraffin

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-07261

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

