Product Datasheet

Cytokeratin 20 Antibody (KRT20/1991) NBP3-07285-100ug

Unit Size: 100 ug Store at 4C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-07285

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-07285




NBP3-07285-100ug

and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array		
Unit Size 100 ug Concentration 0.2 mg/ml Storage Store at 4C. Clonality Monoclonal Clone KRT20/1991 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 2000g/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA 8.0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantity expressed in goblet cells and entercoytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Western Blot. Immunohistochemistry-Paraffin, Protein Array Western Blot. 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 min at RT. Staining of formalin-fixed tissues sections in 10mM Tris with 1 mM EDTA, pH 9.0, for 45 min at 950 Collowed by cooling at RT for 20 minutes.	Cytokeratin 20 Antibody (KRT20/1991)	
Concentration 0.2 mg/ml Storage Store at 4C. Clonality Monoclonal Clone KRT20/1991 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal canner. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and billiary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
Storage Store at 4C. Clonality Monoclonal Clone KRT20/1991 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet call and enterocytes of the qastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Application Notes Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2 ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug
Clone KRT20/1991 Preservative 0.05% Sodium Azide Isotype Ig61 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200g/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, purple and 195C followed by cooling at RT for 20 minutes.	Concentration	0.2 mg/ml
Clone KRT20/1991 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 4kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pt 190, for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at 4C.
Preservative 19G1 Kappa 19G	Clonality	Monoclonal
IgG1 Kappa Purity	Clone	KRT20/1991
Purity Protein A or G purified Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Application Notes Western Blot, Immunohistochemistry-Paraffin, Protein Array Mestern Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating dissue sections in 10mM Tris with 1mM EDTA, pH >0, 0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer 10 mM PBS with 0.05% BSA Target Molecular Weight 46 kDa Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cell and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Application Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG1 Kappa
Target Molecular Weight Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Product Description Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS with 0.05% BSA
Description 200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G. Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	46 kDa
Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT, Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Host Mouse Gene ID 54474 Gene Symbol KRT20 Species Human Marker Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10 mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0 mg/ml. (NBP3-08406) Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to
Gene Symbol KRT20 Species Human Colorectal Epithelial Marker Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Mouse
Species	Gene ID	54474
Marker Colorectal Epithelial Marker This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	KRT20
Specificity/Sensitivity This monoclonal antibody recognizes an intermediate filament protein of 46kDa, identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon, stomach and biliary tract. Breast carcinomas are generally non-reactive. Immunogen Recombinant fragment of human Cytokeratin 20 protein (around aa 196-323) (exact sequence is proprietary) (Uniprot: P35900) Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Colorectal Epithelial Marker
Product Application Details Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Specificity/Sensitivity	identified as cytokeratin 20 (KRT20). KRT is abundantly expressed in goblet cells and enterocytes of the gastrointestinal tract. It is a useful marker of pancreatic and colorectal cancer. KRT20 is expressed under normal, hyperplastic and neoplastic conditions. It has been detected in adenocarcinomas of the colon,
Applications Western Blot, Immunohistochemistry-Paraffin, Protein Array Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	
Recommended Dilutions Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 min at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Western Blot, Immunohistochemistry-Paraffin, Protein Array
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Western Blot 1-2 ug/ml, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array
	Application Notes	formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.




Images


Western Blot: Cytokeratin 20 Antibody (KRT20/1991) [NBP3-07285] - Western Blot Analysis of human HT29 cell lysate using Cytokeratin 20 Mouse Monoclonal Antibody (KRT20/1991).




Immunohistochemistry-Paraffin: Cytokeratin 20 Antibody (KRT20/1991) [NBP3-07285] - Formalin-fixed, paraffin-embedded human Normal Colon stained with Cytokeratin 20 Mouse Monoclonal Antibody (KRT20/1991).



Immunohistochemistry-Paraffin: Cytokeratin 20 Antibody (KRT20/1991) [NBP3-07285] - Formalin-fixed, paraffin-embedded human Colon Carcinoma stained with Cytokeratin 20 Mouse Monoclonal Antibody (KRT20/1991).



Protein Array: Cytokeratin 20 Antibody (KRT20/1991) [NBP3-07285] - Analysis of Protein Array containing more than 19,000 full-length human proteins using Cytokeratin 20 Mouse Monoclonal Antibody (KRT20/1991).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-07285-100ug

HAF007 Goat anti-Mouse IgG Secondary Antibody [HRP]

NB720-B Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]

NBP1-43319-0.5mg Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)
NBP2-23165 Recombinant Human Cytokeratin 20 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-07285

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

