# **Product Datasheet**

## TdT Antibody (DNTT/4506R) NBP3-07684-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-07684

Updated 7/16/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-07684



## NBP3-07684-100ug

TdT Antibody (DNTT/4506R)

· · · · · · · · · · · · · · · · · · ·	
Product Information	
Unit Size	100 ug
Concentration	0.2 mg/ml
Storage	Store at 4C.
Clonality	Monoclonal
Clone	DNTT/4506R
Preservative	0.05% Sodium Azide
Isotype	IgG Kappa
Purity	Protein A or G purified
Buffer	10 mM PBS with 0.05% BSA
Target Molecular Weight	58 kDa
Product Description	
Description	<ul> <li>200ug/ml of antibody purified from Bioreactor Concentrate by Protein A or G.</li> <li>Prepared in 10 mM PBS with 0.05% BSA &amp; 0.05% azide. Also available</li> <li>WITHOUT BSA &amp; azide at 1.0 mg/ml. (NBP3-08564)</li> <li>Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.</li> </ul>
Host	Rabbit
Gene ID	1791
Gene Symbol	DNTT
Species	Human
Marker	Acute Lymphoblastic Leukemia Marker
Immunogen	Synthetic peptide corresponding to residues within aa1-100 of TdT was used as an immunogen. (Uniprot: P04053)
Product Application Details	
Applications	Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunohistochemistry-Paraffin 1-2 ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

www.novusbio.com




#### Images

Immunohistochemistry-Paraffin: TdT Antibody (DNTT/4506R) [NBP3-07684] - IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using TdT antibody (DNTT/4506R) at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2 : HRP-polymer, 30min. DAB, 5min.

Immunohistochemistry-Paraffin: TdT Antibody (DNTT/4506R) [NBP3-07684] - Formalin-fixed, paraffin-embedded human thymus stained with TdT Recombinant Rabbit Monoclonal Antibody (DNTT/4506R).

Immunohistochemistry-Paraffin: TdT Antibody (DNTT/4506R) [NBP3-07684] - Formalin-fixed, paraffin-embedded human thymus stained with TdT Recombinant Rabbit Monoclonal Antibody (DNTT/4506R).







### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-07684-100ug

210-TA-005	TNF-alpha [Unconjugated]
H00001791-Q01-10ug	Recombinant Human TdT GST (N-Term) Protein
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-07684

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

