Product Datasheet

Nidogen-1/Entactin Antibody (ELM1) [Janelia Fluor® 585] NBP3-08237JF585

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08237JF585

Updated 12/12/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08237JF585



NBP3-08237JF585

Nidogen-1/Entactin Antibody (ELM1) [Janelia Fluor® 585]

INICOGET - 1/ETITACTITI ATTRIBUCTY (ELIVIT) [Janetia Fidol & 300]	
Product Information	
Unit Size	0.1 ml
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at 4C in the dark.
Clonality	Monoclonal
Clone	ELM1
Preservative	0.05% Sodium Azide
Isotype	IgG2a Kappa
Conjugate	Janelia Fluor 585
Purity	Protein A or G purified
Buffer	50mM Sodium Borate
Product Description	
Host	Rat
Gene ID	4811
Gene Symbol	NID1
Species	Mouse, Guinea Pig
Reactivity Notes	Does not react with Human.
Specificity/Sensitivity	It reacts with the intact (150kDa) nidogen as well as its proteolytic fragments of 135kDa, 110kDa, 100kDa, and 80kDa. This monoclonal antibody does not cross-react with other basement membrane components or fibronectin. Entactin/Nidogen is a ubiquitous multi-domain basement membrane protein. Basement membranes are the earliest extracellular matrices produced during embryogenesis. They are synthesized and incorporated into the supramolecular architecture of several components, including laminins, Collagen IV, Nidogen and proteoglycans. Nidogen/Entactin, a sulfated glycoprotein, acts as a link between the extracellular matrix molecules Laminin 1 and Collagen Type IV, and thereby participates in the assembly of basement membranes. Nidogen is synthesized and secreted in primary and established mesenchymal peritubular cells and myoepithelial cells, and it affects adhesion of peritubular cells in an autocrine manner. Nidogen is expressed during embryonic and fetal development exclusively in fully developed basement membranes of the ectoderm and is not expressed in the developing endodermal basement membrane or in membranes disrupted during mesoderm formation.
Immunogen	Murine EHS laminin preparation
Notes	Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.
Product Application Details	
Applications	ELISA
Recommended Dilutions	ELISA



Optimal dilution of this antibody should be experimentally determined.

Application Notes



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08237JF585

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

