Product Datasheet

TFF1/pS2 Antibody (TFF1/2969R) - Azide and BSA Free NBP3-08252

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08252

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08252



NBP3-08252


TFF1/pS2 Antibody (TFF1/2969R) - Azide and BSA Free

trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocring therapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	.,	,
Concentration	Product Information	
Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal	Unit Size	100 ug
Clone TFF1/2969R Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 6.5 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein, pS2 is a trefoli peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoli motif, or P-domain, pS2 is known to form dimers and this dimerization is thought to pta role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus, pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPMCFYPNTIDVPPEECCEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: PO4155) Product Application Details Application Notes Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes	Concentration	1 mg/ml
Clone	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Preservative IgG	Clonality	Monoclonal
Isotype IgG	Clone	TFF1/2969R
Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 6.5 kDa Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey Specifficity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus, pS2 is shown to be localized in normal stormach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocring the trapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Application Notes Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes	Preservative	No Preservative
Buffer 10 mM PBS	Isotype	IgG
Product Description	Purity	Protein A or G purified
Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G, Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrintherapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Application Notes Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Buffer	10 mM PBS
1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Rabbit	Target Molecular Weight	6.5 kDa
Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen-receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocring the rapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEECCEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry-Paraffin 1-2 ug/ml	Product Description	
Gene ID Gene Symbol TFF1 Species Human, Cynomolgus Monkey It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocring the rapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Applications Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	•	Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07343). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to
Species	Host	Rabbit
Specificity/Sensitivity	Gene ID	7031
It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocring therapy. Immunogen	Gene Symbol	TFF1
protein. Its epitope is localized between aa57-84 of human p\$2 protein. p\$2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. p\$2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for p\$2. Staining is cytoplasmic, often with localization to the Golgi apparatus. p\$2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that p\$2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine therapy. Immunogen Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/p\$2 (Uniprot: P04155) Product Application Details Applications Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin Immunohistochemistry-Paraffin Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Species	Human, Cynomolgus Monkey
CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155) Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Specificity/Sensitivity	protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine
Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Immunogen	CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of
Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Product Application Details	
Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Applications	Immunohistochemistry-Paraffin
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml
EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Application Notes	



Images

Immunohistochemistry-Paraffin: TFF1/pS2 Antibody (TFF1/2969R) - Azide and BSA Free [NBP3-08252] - Formalin-fixed, paraffin-embedded human Stomach Carcinoma stained with TFF1/pS2 Rabbit Recombinant Monoclonal Antibody (TFF1/2969R).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08252

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-35042-5ug Recombinant Human TFF1/pS2 Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08252

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

