## **Product Datasheet**

# TFF1/pS2 Antibody (TFF1/2969R) [Alexa Fluor® 488] NBP3-08252AF488

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08252AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08252AF488



### NBP3-08252AF488

TFF1/pS2 Antibody (TFF1/2969R) [Alexa Fluor® 488]

| Storage Store at 4C in the dark.  Clonality Monoclonal  Clone TFF1/2969R  Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain, pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive   | TTT Trpo2 Antibody (TTT Tr29091X) [Alexa Fluor® 400] |   |  |
|--|--|---|--|
| Concentration  Please see the vial label for concentration. If unlisted please contact technical services.  Storage  Store at 4C in the dark.  Clonality  Monoclonal  Clone  TFF1/2969R  Preservative  0.05% Sodium Azide  Isotype  IgG  Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Rabbit  Gene ID  7031  Gene Symbol  TFF1  Species  Human, Cynomolgus Monkey  Specificity/Sensitivity  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive | Product Information                                  |   |  |
| Storage Store at 4C in the dark.  Clonality Monoclonal  Clone TFF1/2969R  Preservative 0.05% Sodium Azide  Isotype IgG  Conjugate Alexa Fluor 488  Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 ir trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain, pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positiv   | Jnit Size  | 100 ul  |  |
| Clonality Monoclonal Clone TFF1/2969R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive   | Concentration  | Please see the vial label for concentration. If unlisted please contact technical services.   |  |
| Clone TFF1/2969R  Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 488  Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 7031 Gene Symbol TFF1 Species Human, Cynomolgus Monkey  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive  | Storage  | Store at 4C in the dark.  |  |
| Preservative  Isotype  IgG  Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  Buffer  50mM Sodium Borate  Product Description  Host  Rabbit  Gene ID  7031  Gene Symbol  TFF1  Species  Human, Cynomolgus Monkey  Specificity/Sensitivity  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptides. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are position.   | Clonality  | Monoclonal  |  |
| Isotype Conjugate Alexa Fluor 488  Purity Protein A or G purified Buffer 50mM Sodium Borate  Product Description Host Rabbit Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive  | Clone  | TFF1/2969R  |  |
| Conjugate  Alexa Fluor 488  Purity  Protein A or G purified  50mM Sodium Borate  Product Description  Host  Rabbit  Gene ID  7031  Gene Symbol  TFF1  Species  Human, Cynomolgus Monkey  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 it trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive  | Preservative   | 0.05% Sodium Azide  |  |
| Purity Protein A or G purified  Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.   | sotype   | IgG   |  |
| Buffer 50mM Sodium Borate  Product Description  Host Rabbit  Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive  | Conjugate  | Alexa Fluor 488   |  |
| Product Description  Host Rabbit  Gene ID 7031  Gene Symbol TFF1  Species Human, Cynomolgus Monkey  Specificity/Sensitivity It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive   | Purity   | Protein A or G purified   |  |
| Host  Gene ID  7031  Gene Symbol  TFF1  Species  Human, Cynomolgus Monkey  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive  | 3uffer   | 50mM Sodium Borate  |  |
| Gene Symbol TFF1 Species Human, Cynomolgus Monkey  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.   | Product Description                                  |   |  |
| Gene Symbol  TFF1  Species  Human, Cynomolgus Monkey  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.  | lost   | Rabbit  |  |
| Specificity/Sensitivity  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.   | Gene ID  | 7031  |  |
| Specificity/Sensitivity  It recognizes a polypeptide of 6.5kDa, identified as pS2 estrogen-regulated protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.   | 3ene Symbol  | TFF1  |  |
| protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive.  | Species  | Human, Cynomolgus Monkey  |  |
| pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-  | Specificity/Sensitivity                              | protein. Its epitope is localized between aa57-84 of human pS2 protein. pS2 is a trefoil peptide. Trefoil peptides are protease resistant molecules secreted throughout the gut that play a role in mucosal healing. These peptides contain three intra-chain disulfide bonds, forming the trefoil motif, or P-domain. pS2 is known to form dimers and this dimerization is thought to play a role in its protective and healing properties. About 60% of breast carcinomas are positive for pS2. Staining is cytoplasmic, often with localization to the Golgi apparatus. pS2 is shown to be localized in normal stomach mucosa, gastric fluid, goblet cells in the colon and small intestine, and in ulcerations of the gastrointestinal tract. Several studies have shown that pS2 is primarily expressed in estrogen receptor-positive breast tumors and it may define a subset of estrogen-dependent tumors that displays an increased likelihood of response to endocrine |  |
| Synthetic peptide of 28 amino acid residues corresponding to CFDDTVRGVPWCFYPNTIDVPPEECCEF (aa57-84) from the C-terminus of human TFF1/pS2 (Uniprot: P04155)  | mmunogen   | CFDDTVRGVPWCFYPNTIDVPPEEECEF (aa57-84) from the C-terminus of   |  |



#### Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

| <b>Product Application Details</b> |  |
|------------------------------------|--|
| Applications                       | Immunohistochemistry-Paraffin  |
| Recommended Dilutions              | Immunohistochemistry-Paraffin  |
| Application Notes                  | Optimal dilution of this antibody should be experimentally determined. |





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### **Products Related to NBP3-08252AF488**

NBP2-24982 Rabbit IgG Isotype Control [Alexa Fluor® 488]

NBP2-35042-5ug Recombinant Human TFF1/pS2 Protein

236-EG-200 EGF [Unconjugated]
DY5237 TFF1/pS2 [Biotin]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08252AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

