# **Product Datasheet**

## Integrin beta 3/CD61 Antibody (ITGB3/2166R) [Alexa Fluor® 532] NBP3-08291AF532

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08291AF532

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-08291AF532



## NBP3-08291AF532

Integrin beta 3/CD61 Antibody (ITGB3/2166R) [Alexa Fluor® 532]

U		
Product	Information	

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneITGB3/2166RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer3690Gene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpeciesReact swith human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41/CD61 complex apears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for on willebrand factor, soluble fibrinogen, fibronectin, witronectin and aggregation. The CD51/CD61 is implicated in unror metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes arted ration cole is proprietary) (Uniprot: P05106)	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneITGB3/2166RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpeciesSects with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61 complex sia a receptor for von Wilebrand factor, soluble fibringen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adnoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Unit Size	100 ul	
ClonalityMonoclonalCloneITGB3/2166RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor bet a chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex a central role in platelet activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and agregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Concentration		
CloneITGB3/2166RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Clonality	Monoclonal	
IsotypeIgGConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Clone	ITGB3/2166R	
ConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Isotype	IgG	
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex are receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Conjugate	Alexa Fluor 532	
Product DescriptionHostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Purity	Protein A or G purified	
HostRabbitGene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Buffer	50mM Sodium Borate	
Gene ID3690Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Product Description		
Gene SymbolITGB3SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Host	Rabbit	
SpeciesHumanMarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Gene ID	3690	
MarkerPlatelet MarkerSpecificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Gene Symbol	ITGB3	
Specificity/SensitivityReacts with human integrin beta3 (GPIIIa, vitronectin receptor beta chain). It associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Species	Human	
associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61). The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.ImmunogenRecombinant human Integrin beta 3/CD61 protein fragment (aa 385-490) (exact	Marker	Platelet Marker	
	Specificity/Sensitivity	associates with the IIb-chain (CD41) to form the GpIIb/GpIIIa complex (CD41/CD61).The CD41/CD61 complex appears early in megakaryocyte maturation. The activated CD41/CD61 complex is a receptor for von Willebrand factor, soluble fibrinogen, fibronectin, vitronectin and thrombospondin. It plays a central role in platelet activation and aggregation. The CD51/CD61 is implicated in tumor metastasis and adenoviral infection. The antibody detects platelets in smears of blood and bone marrow, as well as megakaryocytes in frozen sections and cell smears. The antibody is useful for classification of megakaryoblastic leukemia.	
	Immunogen		

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.	
Product Application Details		
Applications	Immunohistochemistry-Paraffin	
<b>Recommended Dilutions</b>	Immunohistochemistry-Paraffin	
Application Notes	Optimal dilution of this antibody should be experimentally determined.	

Notes





## Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

## Products Related to NBP3-08291AF532

NBL1-12073	Integrin beta 3/CD61 Overexpression Lysate
210-TA-005	TNF-alpha [Unconjugated]
NBP1-83453PEP	Integrin beta 3/CD61 Recombinant Protein Antigen
NBP2-24891AF532	Rabbit IgG Isotype Control [Alexa Fluor® 532]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08291AF532

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

