Product Datasheet

PAX8 Antibody (PAX8/3688R) [Alexa Fluor® 647] NBP3-08295AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08295AF647

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08295AF647



NBP3-08295AF647

PAX8 Antibody (PAX8/3688R) [Alexa Fluor® 647]

Product Information Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PAX8/3688R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 7849 Gene Symbol PAX8 Specificity/Sensitivity Renal Cell Marker Specificity/Sensitivity Recognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid adenomas. PAX-8 is expressed in the thyroid adenomas. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage ovarian serous, epithelial cells. PAX-8 is expressed in a high percentage ovarian serous, epithelial celle carcin	, L	•
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Stor at 4C in the dark. Clonality Monoclonal Clone PAX8/3688R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Rabbit Gene ID 7849 Gene Symbol PAX8 Species Human Marker Renal Cell Marker Specificity/Sensitivity Recognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid folicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid folicular ceinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian serous, endometrioid, and clear cell carcinomas, BAX-8 is expressed in the thyroid (and clear cell carcinomas, PAX-8 is expressed in the thyroid of and clear cell carcinomas, PAX-8 attibody may be used as an additional immunohistochemical marker for renal epithelial attimosi.	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePAX8/3688RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription and atypical actionaus. PAX-8 is expressed in the thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cells. PAX-8 is expressed in a high percentage of ovarian servus, endometrioid, and clear cell carcinomas, PAX-8 but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian servus, endometrioid, and clear cell carcinomas, PAX-8 antibody may be use	Unit Size	100 ul
ClonalityMonoclonalClonePAX8/3688RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerReal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cerli development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations and atypical thyroid acter or normal, ovarian serous, endometricid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarian inclusion cysts, but not normal ovarian serous, endometricid, and clear cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next-8 expression is reported in renal tubules as well as renal cell carcinoma, next	Concentration	
ClonePAX8/3688RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid specific and sociated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid specific and sociated acrinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunchistochemical marker for renal epithelial tumors.	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thryroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, pat A expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular cell development and expression is reported in renal tubules as well as renal cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, pAX-8 and seriores on seriors. PAX-8 and seriors and additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Clone	PAX8/3688R
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid specific genes. Mutations in this gene have been associated with thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Conjugate	Alexa Fluor 647
Product DescriptionHostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas, non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, pAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial turrors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Purity	Protein A or G purified
HostRabbitGene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Buffer	50mM Sodium Borate
Gene ID7849Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas. but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Product Description	
Gene SymbolPAX8SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinoma. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Host	Rabbit
SpeciesHumanMarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Gene ID	7849
MarkerRenal Cell MarkerSpecificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Gene Symbol	PAX8
Specificity/SensitivityRecognizes a protein of 62kDa, identified as PAX8. It is a member of the paired box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors.ImmunogenSynthetic peptide corresponding to the C-terminus of human PAX8 protein	Species	Human
 box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial tumors. Immunogen 	Marker	Renal Cell Marker
	Specificity/Sensitivity	box (PAX) family of transcription factors. This nuclear protein is involved in thyroid follicular cell development and expression of thyroid-specific genes. Mutations in this gene have been associated with thyroid dysgenesis, thyroid follicular carcinomas, and atypical thyroid adenomas. PAX-8 is expressed in the thyroid (and associated carcinomas), non-ciliated mucosal cells of the fallopian tubes, and simple ovarian inclusion cysts, but not normal ovarian surface epithelial cells. PAX-8 is expressed in a high percentage of ovarian serous, endometrioid, and clear cell carcinomas, but only rarely in primary ovarian mucinous adenocarcinomas. PAX-8 expression is reported in renal tubules as well as renal cell carcinoma, nephroblastoma, and seminoma. PAX-8 antibody may be used as an additional immunohistochemical marker for renal epithelial
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08295AF647

NBP2-24891AF647	Rabbit IgG Isotype Control [Alexa Fluor® 647]
NBP2-51907-0.05mg	Recombinant Human PAX8 His Protein
288-TPN-025/CF	Thrombopoietin/THPO [Unconjugated]
NBL1-14132	PAX8 Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08295AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

