## **Product Datasheet**

### Bcl-6 Antibody (BCL6/2497R) [Alexa Fluor® 647] NBP3-08313AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08313AF647

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08313AF647



#### NBP3-08313AF647

Bcl-6 Antibody (BCL6/2497R) [Alexa Fluor® 647]

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneBCL6/2497RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604Gene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large lecell lymphomas (DLBCL), where CD10, bcl-6 and MUM1//RF4 are used to identify germinal center and activated B-cell lymphoma are local bodyin lymphoma (DLBCL), where cD10, bcl-6 and MUM1//RF4 are used to identify germinal center and activated B-cell lymphoma are local bodyin lymphoma are bcl-6 negative whereas the large (LH) cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 arrety stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecominant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary) (Uniprot: P41182)		
services.StorageStor at 4C in the dark.ClonalityMonoclonalCloneBCL6/2497RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604Gene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphoma (L)_PML), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Unit Size	100 ul
ClonalityMonoclonalCloneBCL6/2497RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description604HostRabbitGene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFolicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as BcI-6. Antibody to bcI-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma are usually negative. (2) BcI-6 is an important prognostic marker in diffuse large B-cell lymphoma (DLBCL), where CD10, bcI-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell hymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of Classical Hodgkin lymphoma.ImmunogenRecombinant human bcI-6 protein fragment (aa256-389) (exact sequence is	Concentration	
CloneBCL6/2497RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Storage	Store at 4C in the dark.
Preservative       0.05% Sodium Azide         Isotype       IgG         Conjugate       Alexa Fluor 647         Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description       Host         Host       Rabbit         Gene ID       604         Gene Symbol       BCL6         Species       Human, Canine         Marker       Follicular Lymphoma Marker         Specificity/Sensitivity       Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma.         Immunogen       Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Clonality	Monoclonal
Isotype       IgG         Conjugate       Alexa Fluor 647         Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description       Host         Host       Rabbit         Gene ID       604         Gene Symbol       BCL6         Species       Human, Canine         Marker       Follicular Lymphoma Marker         Specificity/Sensitivity       Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.         Immunogen       Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Clone	BCL6/2497R
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/RF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Preservative	0.05% Sodium Azide
Purity       Protein A or G purified         Buffer       50mM Sodium Borate         Product Description          Host       Rabbit         Gene ID       604         Gene Symbol       BCL6         Species       Human, Canine         Marker       Follicular Lymphoma Marker         Specificity/Sensitivity       Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.         Immunogen       Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Conjugate	Alexa Fluor 647
Product Description         Host       Rabbit         Gene ID       604         Gene Symbol       BCL6         Species       Human, Canine         Marker       Follicular Lymphoma Marker         Specificity/Sensitivity       Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.         Immunogen       Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Purity	Protein A or G purified
HostRabbitGene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Buffer	50mM Sodium Borate
Gene ID604Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Product Description	
Gene SymbolBCL6SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Host	Rabbit
SpeciesHuman, CanineMarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Gene ID	604
MarkerFollicular Lymphoma MarkerSpecificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Gene Symbol	BCL6
Specificity/SensitivityRecognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.ImmunogenRecombinant human bcl-6 protein fragment (aa256-389) (exact sequence is	Species	Human, Canine
<ul> <li>helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.</li> <li>Immunogen</li> </ul>	Marker	Follicular Lymphoma Marker
	Specificity/Sensitivity	helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma.
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-08313AF647

NBP2-24891AF647	Rabbit IgG Isotype Control [Alexa Fluor® 647]
H00000604-Q01-10ug	Recombinant Human Bcl-6 GST (N-Term) Protein
594-ML-010	IL-21 [Unconjugated]
NBL1-07951	Bcl-6 Overexpression Lysate

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08313AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

