Product Datasheet

Golgi Complex Antibody (GLG1/2829R) [Alexa Fluor® 647] NBP3-08396AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08396AF647

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08396AF647



NBP3-08396AF647

Golgi Complex Antibody (GLG1/2829R) [Alexa Fluor® 647]

Product Information

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneGLG1/2829RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHoman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in cells or tissue preparations and can be used to stain the Golgi complex in cells or tissue preparations and can be used as a Golgi marker for Human cells in xenographic model research. It reacts specifically with human cells or use within the cells. This monoclonal antibody rescipation or the Golgi complex in cells or tissue preparations and can be used as a Golgi marker for subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody rescipation of the Golgi zone in normal and malignant cells. This monoclonal antibody rescipation of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Sodigi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell to rescipate sit or digi apparatus is to for for Golgi arparents the colgi are present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules passing though the	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone GLG1/2829R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Rabbit Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xeenographic model research. It reacts specifically with than cells. The Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sace schown as citernae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules served spherical vescides that have bud	Unit Size	100 ul
ClonalMonoclonalCloneGLG1/2829RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi apparatus is an organelle present. It reacts specifically with human cells in xenographic model research. It reacts specifically with human cells in the cells. This monoclonal antibody is an excellent marker for Human cells in scengaphic model research. It reacts specifically with human cells in the cell in access and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sack shown as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules spherical vesites that have budded off of the Golgi are present surrounding the main cisternae.	Concentration	
CloneGLG1/2829RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with numan cells in xenographic model research. It reacts specifically with numan cells in xenographic model research. It reacts specifically with human cells for sectors or use within the cell. The Golgi and parkage macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Host Rabbit Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used to stain the Golgi complex in cells or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in senographic model research. It reacts specifically with human cells in senographic model research. It reacts specifically with human cells in senographic model research. It reacts specifically with human cells in senographic model research. It reacts specifically with human cells in senographic model research. It reacts the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Clone	GLG1/2829R
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in the cell. The Golgi is made up of a stack of flattned, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively molify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Isotype	IgG
Buffer 50mM Sodium Borate Product Description Rabbit Host Rabbit Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Conjugate	Alexa Fluor 647
Product Description Host Rabbit Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Marker for Human Cells Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in cell or tissue preparations and can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Purity	Protein A or G purified
HostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Buffer	50mM Sodium Borate
SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane- bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Product Description	
Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane- bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Host	Rabbit
MarkerMarker for Human CellsSpecificity/SensitivityThis monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane- bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Species	Human, Mouse (Negative), Rat (Negative)
Specificity/Sensitivity This monoclonal antibody recognizes an antigen associated with the Golgi complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane-bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Reactivity Notes	Does not react with Mouse or Rat.
complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane- bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded off of the Golgi are present surrounding the main cisternae.	Marker	Marker for Human Cells
Immunogen Recombinant full-length human Golgi Complex protein (Uniprot: Q92896)	Specificity/Sensitivity	complex in human cells only. It can be used to stain the Golgi complex in cell or tissue preparations and can be used as a Golgi marker in subcellular fractions. It produces a diffuse staining pattern of the Golgi zone in normal and malignant cells. This monoclonal antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells. The Golgi apparatus is an organelle present in all eukaryotic cells that forms a part of the endomembrane system. The primary function of the Golgi apparatus is to process and package macromolecules synthesized by the cell for exocytosis or use within the cell. The Golgi is made up of a stack of flattened, membrane- bound sacs known as cisternae, with three functional regions: the cis face, medial region and trans face. Each region consists of various enzymes that selectively modify the macromolecules passing though them, depending on where they are destined to reside. Several spherical vesicles that have budded
	Immunogen	Recombinant full-length human Golgi Complex protein (Uniprot: Q92896)

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08396AF647

NBP2-24891AF647

Rabbit IgG Isotype Control [Alexa Fluor® 647]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08396AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

