Product Datasheet

Carbonic Anhydrase IX/CA9 Antibody (CA9/2993R) [Alexa Fluor® 700] NBP3-08407AF700

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08407AF700

Updated 9/11/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-08407AF700



NBP3-08407AF700

Carbonic Anhydrase IX/CA9 Antibody (CA9/2993R) [Alexa Fluor® 700]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCA9/2993RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID768Gene SymbolCA9Specificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CA1X/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivi		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCA9/2993RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID768Gene SymbolCA9Specificity/SensitivityRecognizes a glycoprotein of -200kDa, identified as carbonic anhydrase IX (CAIX/g200). Its epitope resides in the carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group solution and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Byomans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is sequresed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneCA9/2993RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/g200). Its epitope resides in the carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group solicized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, g200 is sequesed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Unit Size	100 ul
ClonalityMonoclonalCloneCA9/293RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp20 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal nd 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Concentration	•
CloneCA9/2993RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate domain of gp200. It shows or significant cross-reactivity with other carbohydrate domain of gp200. It shows or significant cross-reactivity with other carbohydrate domain of gp200. It shows or significant cross-reactivity with ot	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 700 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 768 Gene Symbol CA9 Species Human, Equine Species Human, Equine Specificity/Sensitivity Recognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation. Immunogen Recombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Clone	CA9/2993R
ConjugateAlexa Fluor 700PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Conjugate	Alexa Fluor 700
Product DescriptionHostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Purity	Protein A or G purified
HostRabbitGene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Buffer	50mM Sodium Borate
Gene ID768Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Product Description	
Gene SymbolCA9SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Host	Rabbit
SpeciesHuman, EquineSpecificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Gene ID	768
Specificity/SensitivityRecognizes a glycoprotein of ~200kDa, identified as carbonic anhydrase IX (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation.ImmunogenRecombinant full-length human Carbonic Anhydrase IX/CA9 protein (Uniprot:	Gene Symbol	CA9
 (CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially those showing tubular differentiation. Immunogen 	Species	Human, Equine
	Specificity/Sensitivity	(CAIX/gp200). Its epitope resides in the carbohydrate domain of gp200. It shows no significant cross-reactivity with other carbohydrate determinants, such as the Lewis blood group antigens, epithelial membrane antigen, HMFG, and AB blood group antigens. In normal kidney, gp200 is localized along the brush border of the pars convoluta and pars recta segments of the proximal tubule, as well as focally along the luminal surface of Bowmans capsule adjoining the outgoing proximal tubule. Reportedly, gp200 is expressed by 93% of primary and 84% of metastatic renal cell carcinomas. This monoclonal antibody may be useful in the investigations of carcinomas of proximal nephrogenic differentiation especially
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on
	demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08407AF700

DCA900	Carbonic Anhydrase IX/CA9 [HRP]
210-TA-005	TNF-alpha [Unconjugated]
NB100-417PEP	Carbonic Anhydrase IX/CA9 Antibody Blocking Peptide
NBP2-24891AF700	Rabbit IgG Isotype Control [Alexa Fluor® 700]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08407AF700

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

