Product Datasheet

Desmoglein-3 Antibody (DSG3/2796) [Alexa Fluor® 594] NBP3-08410AF594

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08410AF594

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08410AF594



NBP3-08410AF594

Desmoglein-3 Antibody (DSG3/2796) [Alexa Fluor® 594]

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneDSG3/2796Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionIso30Gene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquanous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). The second and the protein component of desmosomes in vertebrate epithelial specificity/Sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact sequence is proprietary) (Uniprot: P329	Froduct mormation	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone DSG3/2796 Preservative 0.05% Sodium Azide Isotype IgG2b Kappa Conjugate Alexa Fluor 594 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Hose Host Mouse Gene ID 1830 Gene Symbol DSG3 Species Human Marker Squamous Cell Marker Specificity/Sensitivity Recognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-anily. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DS	Unit Size	100 ul
ClonalityMonoclonalCloneDSG3/2796Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHouseHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Concentration	•
CloneDSG3/2796Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 in 2003). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). Therefore, DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Clonality	Monoclonal
IsotypeIgG2b KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Clone	DSG3/2796
ConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmebrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Isotype	IgG2b Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Conjugate	Alexa Fluor 594
Product DescriptionHostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Purity	Protein A or G purified
HostMouseGene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Buffer	50mM Sodium Borate
Gene ID1830Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Product Description	
Gene SymbolDSG3SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Host	Mouse
SpeciesHumanMarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Gene ID	1830
MarkerSquamous Cell MarkerSpecificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Gene Symbol	DSG3
Specificity/SensitivityRecognizes a protein of 130kDa, identified as Desmoglein-3 (DSG3). This monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis.ImmunogenRecombinant fragment (around aa 379-491) human Desmoglein-3 protein (exact	Species	Human
 monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate a poor prognosis. Immunogen 	Marker	Squamous Cell Marker
	Specificity/Sensitivity	monoclonal antibody is highly specific to Desmoglein-3 and does not cross-react with other members of the Desmoglein-family. DSG3 is a calcium-binding transmembrane glycoprotein component of desmosomes in vertebrate epithelial cells. Research has shown that DSG3 has a very high sensitivity (80%) and specificity (100%) in recognizing squamous cell carcinoma (SqCC). Therefore, DSG3 is considered a very important marker for lung SqCC and can be a useful ancillary marker to separate SqCC from other subtypes of lung cancer. Moreover, studies have shown that DSG3 expression in lung SqCC may indicate
	Immunogen	



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, ELISA, Protein Array
Recommended Dilutions	Western Blot, ELISA, Protein Array
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08410AF594

NBP1-43317AF594Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor®
594]NBP2-54744PEPDesmoglein-3 Recombinant Protein Antigen6507-IL-010/CFIL-4 [Unconjugated]11167-DM-050Desmoglein-3 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08410AF594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

