Product Datasheet

dsDNA Antibody (rDSD/4565) [Biotin] NBP3-08466B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08466B

Updated 3/28/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08466B



NBP3-08466B

dsDNA Antibody (rDSD/4565) [Biotin]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone rDSD/4565 Preservative 0.05% Sodium Azide Isotype IgG2a Kappa Conjugate Biotin Purity Protein A purified Buffer PBS Product Description Mouse Host Mouse Species Human Species Human Species Human Specificity/Sensitivity This monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nucleu in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleative. The onormer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by sydrogen b	ConcentrationPlease as servicesStorageStore atClonalityMonocloClonerDSD/45Preservative0.05% SIsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionHumanMarkerNuclearSpeciesHumanMarkerSubcelluuseful in preparationproduce and on be u produce malignan material monome one of for together two come by hydrome	C in the dark. nal 55 odium Azide ppa
services.StorageStore at 4C in the dark.ClonalityMonoclonalClonerDSD/4565Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateBiotinPurityProtein A purifiedBufferBBSProduct DescriptionMouseMarkerNuclear MarkerSpeciesHumanSpeciesThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. This monoclonal antibody recognizes the double stranded DNA in human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. These morkers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody recognizes the induced of stain the nuclei in cell or tissue preparations and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic water and ys aphosphoristed 2-deoxyribose N-glycosidically linked to gether by aphosphoristed 2-deoxyribose N-glycosidically lin	StorageStore atStorageStore atClonalityMonocloClonerDSD/45Preservative0.05% SIsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionHumanMarkerNuclearSpeciesHumanMarkerSubcelluuseful in preparat human or can be u produce malignai material monome one of for together two com by hydro	C in the dark. nal 55 odium Azide ppa
ClonalityMonoclonalClonerDSD/4565Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateBiotinPurityProtein A purifiedBufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nucleus of normal and matignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked to 	ClonalityMonocloClonerDSD/45Preservative0.05% SIsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionHumanHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis monome one of for together two com by hydro	al 55 odium Azide ppa
ClonerDSD/4565Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateBiotinPurityProtein A purifiedBufferPBSProduct DescriptionHoasHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. These markers may be useful and cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyriboxe N-glycosidically linked to one of four bases, adenine, cytosine, guanine or furkine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin	ClonerDSD/45Preservative0.05% SIsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionHumanHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mosubcellu useful im preparation human of can be u produce malignation material monome one of for 	5 odium Azide ppa
Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateBiotinPurityProtein A purifiedBufferPBSProduct DescriptionMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells. These markers may be useful in identification of these organelles in cells. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nuclei used of stain the nuclei in cell or tissue preparations and 	Preservative0.05% SIsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mosubcellu useful in preparation human of can be u produce malignar material monome one of for together two come by hydrome	odium Azide ppa
IsotypeIgG2a KappaConjugateBiotinPurityProtein A purifiedBufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in 	IsotypeIgG2a KConjugateBiotinPurityProtein ABufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mosubcellu useful in preparation human of can be u produce malignation material monome one of for together two come	рра
ConjugateBiotinPurityProtein A purifiedBufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkit's cellsProduct Application DetailisELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	ConjugateBiotinPurityProtein /BufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mosubcellu useful in preparation human of can be u produce malignation material monome one of for together two come	
PurityProtein A purifiedPurityProtein A purifiedBufferPBSProduct DescriptionHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. T	PurityProtein ABufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mo subcellu useful in preparat human oc can be u produce malignat material monome one of for together two com by hydro	purified
BufferPBSProduct DescriptionMouseHostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	BufferPBSProduct DescriptionHostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mo subcellu useful in preparat human of can be u produce malignat material monome one of for together two com by hydro	purified
Product Description Host Mouse Species Human Marker Nuclear Marker Specificity/Sensitivity This monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs. Immunogen Nuclei of Burkitt's cells Product Application Details ELISA, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin	Product Description Host Mouse Species Human Marker Nuclear Specificity/Sensitivity This mosubcellur useful im preparate human of can be uproduce malignation material monome one of for together two com by hydro	
HostMouseSpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleous of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	HostMouseSpeciesHumanMarkerNuclearSpecificity/SensitivityThis mo subcellu useful in preparat human of can be u produce malignat material monome one of for together two com by hydro	
SpeciesHumanMarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	SpeciesHumanMarkerNuclearSpecificity/SensitivityThis mo subcellu useful in preparat human of can be u produce malignat material monome one of fo together two com by hydro	
MarkerNuclear MarkerSpecificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenKuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	Marker Nuclear Specificity/Sensitivity This mossubcelluuuseful in preparathuman ocan be uproduce malignatimaterial monome one of for together two com by hydrome of the two com by hydrome of two com by hydrome of the two com by hydrome of two	
Specificity/SensitivityThis monoclonal antibody is part of a new panel of reagents, which recognizes subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleus of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	Specificity/Sensitivity This mo subcellu useful in preparat human o can be u produce malignat material monome one of fo together two com by hydro	
subcellular organelles or compartments of human cells. These markers may be useful in identification of these organelles in cells, tissues, and biochemical preparations. This monoclonal antibody recognizes the double stranded DNA in human cells. It can be used to stain the nuclei in cell or tissue preparations and can be used as a nuclear marker in human cells. This monoclonal antibody produces a homogeneous staining pattern in the nucleoi of normal and malignant cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic material of all cells and many viruses and is a polymer of nucleotides. The monomer consists of phosphorylated 2-deoxyribose N-glycosidically linked to one of four bases, adenine, cytosine, guanine or thymine. These are linked together by 3-phosphodiester bridges. In the Watson-Crick double-helix model, two complementary strands are wound in a right-handed helix and held together by hydrogen bonds between complementary base pairs.ImmunogenNuclei of Burkitt's cellsProduct Application DetailsELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	subcellu useful in preparat human o can be u produce malignat material monome one of fo together two com by hydro	<i>l</i> arker
Product Application Details Applications ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin		ar organelles or compartments of human cells. These markers may be dentification of these organelles in cells, tissues, and biochemical ons. This monoclonal antibody recognizes the double stranded DNA in ells. It can be used to stain the nuclei in cell or tissue preparations and sed as a nuclear marker in human cells. This monoclonal antibody a homogeneous staining pattern in the nucleus of normal and t cells. Double Stranded deoxyribonucleic acid (ds DNA) is the genetic of all cells and many viruses and is a polymer of nucleotides. The consists of phosphorylated 2-deoxyribose N-glycosidically linked to ur bases, adenine, cytosine, guanine or thymine. These are linked by 3-phosphodiester bridges. In the Watson-Crick double-helix model, elementary strands are wound in a right-handed helix and held together
Applications ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin	Immunogen Nuclei o	Burkitt's cells
Paraffin	Product Application Details	
Recommended Dilutions ELISA, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry- Paraffin		nmunocytochemistry/ Immunofluorescence, Immunohistochemistry-
Application NotesOptimal dilution of this antibody should be experimentally determined.	Application Notes Optimal	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08466B

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

