Product Datasheet

Mammaglobin A Antibody (MGB/4811R) - Azide and BSA Free NBP3-08488

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08488

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08488

NBP3-08488


Mammaglobin A Antibody (MGB/4811R) - Azide and BSA Free

Product Information Unit Size 100 ug Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone MGB/4811R Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Gene ID 4250 Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin Application Notes Immunohistochemistry-Paraffin Application Notes Immunohistochemistry-Paraffin		
Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone MGB/4811R Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description 10 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Gene ID 4250 Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of tormalin-fixed tissues requires heating tissue sections in 100m Tris with 1rmM	Product Information	
Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone MGB/4811R Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Gene ID 4250 Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunobistochemistry-Paraffin Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details Immunohistochemistry-Paraffin 1-2 ug/ml Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed in at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug
Clonality Monoclonal Clone MGB/4811R Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Rabbit Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details Immunohistochemistry-Paraffin 1-2 ug/ml Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	1 mg/ml
CloneMGB/4811RPreservativeNo PreservativeIsotypeIgGPurityProtein A or G purifiedBuffer10 mM PBSTarget Molecular Weight10.5 kDaProduct Description1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.HostRabbitGene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tis with 1mM EDTA, PI 9, 0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Rabbit Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry-Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description Description Multiple State Antibody with of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Rabbit Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details // Marker Applications // Immunohistochemistry-Paraffin Recommended Dilutions // Immunohistochemistry-Paraffin 1-2 ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM	Clone	MGB/4811R
Purity Protein A or G purified Buffer 10 mM PBS Target Molecular Weight 10.5 kDa Product Description	Preservative	No Preservative
Buffer10 mM PBSTarget Molecular Weight10.5 kDaProduct Description1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.HostRabbitGene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG
Target Molecular Weight10.5 kDaProduct Description1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.HostRabbitGene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-Paraffin Recommended DilutionsApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
Product Description Description 1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C. Host Rabbit Gene ID 4250 Gene Symbol SCGB2A2 Species Human Marker Breast Cancer Marker Immunogen Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10 mM PBS
Description1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.HostRabbitGene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Target Molecular Weight	10.5 kDa
Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300).Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.HostRabbitGene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID4250Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2 ug/mlRecommended DilutionsImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07300). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80
Gene SymbolSCGB2A2SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Rabbit
SpeciesHumanMarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	4250
MarkerBreast Cancer MarkerImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	SCGB2A2
ImmunogenRecombinant full-length human Mammaglobin A protein (Uniprot: Q13296)Product Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
Product Application Details Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Breast Cancer Marker
ApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Recombinant full-length human Mammaglobin A protein (Uniprot: Q13296)
Recommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application NotesImmunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry-Paraffin
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml
	Application Notes	formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.



Images

Immunohistochemistry-Paraffin: Mammaglobin A Antibody (MGB/4811R) - Azide and BSA Free [NBP3-08488] - Formalin-fixed, paraffinembedded human lactating breast stained with Mammaglobin A Recombinant Rabbit Monoclonal Antibody (MGB/4811R).



Immunohistochemistry-Paraffin: Mammaglobin A Antibody (MGB/4811R) - Azide and BSA Free [NBP3-08488] - Formalin-fixed, paraffinembedded human lactating breast stained with Mammaglobin A Recombinant Rabbit Monoclonal Antibody (MGB/4811R).





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08488

anti-Rabbit IgG Secondary Antibody [HRP]
anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
bit IgG Isotype Control
ombinant Human Mammaglobin A GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

