Product Datasheet

Histone H3 [p Ser10] Antibody (PHH3/471R) [Alexa Fluor® 350] NBP3-08511AF350

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08511AF350

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08511AF350



NBP3-08511AF350

Histone H3 [p Ser10] Antibody (PHH3/471R) [Alexa Fluor® 350]

Product Information	Thistorie 113 [p Ser 10] Antibody (FTIT13/47 TK) [Alexa Fluot® 350]	
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PHH3/471R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late 62 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone PHH3/471R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphonistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was ascoitated with increased expression of PHH3 was significantly associated with increased expression of PHH3 was significantly associated with increased expression of PHH3 was subject of tumor pucleration (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Unit Size	100 ul
Clone PHH3/471R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Concentration	
Clone PHH3/471R Preservative 0.05% Sodium Azide Isotype IgG Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Storage	Store at 4C in the dark.
Preservative Sodium Azide	Clonality	Monoclonal
IgG	Clone	PHH3/471R
Conjugate Alexa Fluor 350 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.	Isotype	IgG
Buffer 50mM Sodium Borate	Conjugate	Alexa Fluor 350
Product Description Host Rabbit Gene ID 126961 Gene Symbol H3C14 Species Human Marker Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Purity	Protein A or G purified
Rabbit	Buffer	50mM Sodium Borate
Gene Symbol Gene Symbol H3C14 Species Human Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Product Description	
Gene Symbol H3C14 Species Human Nuclear Marker Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Host	Rabbit
Species Human Nuclear Marker Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Gene ID	126961
Marker Nuclear Marker Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Gene Symbol	H3C14
Specificity/Sensitivity Phosphohistone H3 (PHH3) is a marker specific for cells undergoing mitosis. Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p = 0.041) and tumor necrosis (p = 0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Species	Human
Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis. Immunogen A synthetic peptide corresponding to (ARK-pS-TGGKAPRKQLc) of	Marker	Nuclear Marker
	Specificity/Sensitivity	Serine 10 of Histone H3 is phosphorylated in association with mitotic chromatin condensation in late G2 and M phase of the cell cycle and thus, PHH3 can distinguish mitosis from apoptotic nuclei. The range of percentage PHH3 positive tumor nuclei was from 0.0 to 6.6% (median value 0.8%). Increased expression of PHH3 was significantly associated with tumor thickness (p = 0.031), presence of tumor ulceration (p =0.041) and tumor necrosis (p =0.027), but not with Clarks level of invasion. High levels of PHH3 was associated with increased mitotic count (p = 0.003) and high Ki-67 expression (p = 0.002). For central nervous system tumors, melanoma, soft tissue tumors, GIST, etc., PHH3 mAb is helpful for tumor pathological classification and prognosis.
	Immunogen	



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08511AF350

NBP2-24891AF350 Rabbit IqG Isotype Control [Alexa Fluor® 350]

NB21-1141PEP Histone H3 [Monomethyl Lys18] Antibody Blocking Peptide

H00008362-P01-2ug Recombinant Human H4/d GST (N-Term) Protein

NB21-1251PEP Histone H3 [Monomethyl Lys36] Antibody Blocking Peptide

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08511AF350

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

