Product Datasheet

Collagen IV Antibody (M3F7) [Janelia Fluor® 635] NBP3-08515JF635

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08515JF635

Updated 12/12/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08515JF635



NBP3-08515JF635

Collagen IV Antibody (M3F7) [Janelia Fluor® 635]

Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	conagon it / massay (mor / jean	iona i idolo della
Please see the vial label for concentration. If unlisted please contact technical services.	Product Information	
Storage Store at 4C in the dark. Clonality Monoclonal Clone M3F7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 635 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 1282 Gene Symbol COL4A1 Species Human, Rat, Avian Reactivity Notes Quail Specificity/Sensitivity Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entacturi/niclogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogeneia calviny. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney. Lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen alpha1 (IV)-alpha2(IV) triple helix. The epitope is triple helical domain about 900 from the N-terminus Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Unit Size	0.1 ml
Clone M3F7 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 635 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 1282 Gene Symbol COL4A1 Species Human, Rat, Avian Reactivity Notes Quail Specificity/Sensitivity Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV duse in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/indogen. Arresten, comprising the C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain,	Concentration	·
Clone M3F7 Preservative 0.05% Sodium Azide Isotype 1gG1 Kappa Conjugate Janelia Fluor 635 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 1282 Gene Symbol COL4A1 Species Human, Rat, Avian Reactivity Notes Quail Specificity/Sensitivity Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV tables include use in pancreatic adenocarcinoma and chronic pancreatitis, nephroselerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/hidogen. Arresten, comprising the C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 domain, inhibits angiogenesis and tumor formation. The C-terminal NG1 found in M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen alpha1 (IV)-alpha2(IV) triple helix. The epitope is triple helical domain about 900 from the N-terminus Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Storage	Store at 4C in the dark.
Preservative IgG1 Kappa IgG1 Kappa IgG1 Kappa IgG1 Kappa IgG1 Kappa Janelia Fluor 635 Purity	Clonality	Monoclonal
IgG1 Kappa	Clone	M3F7
Conjugate Janelia Fluor 635 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 1282 Gene Symbol COL4A1 Species Human, Rat, Avian Reactivity Notes Quail Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogeneis and tumor formation. The C-terminal NC1 domain, inhibits angiogeneis and tumor formation. The C-terminal Cell proliferation, migration, and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen slother license from the Howard Hughes Medical Institute, Janelia Research Campus.	Preservative	0.05% Sodium Azide
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 1282 Gene Symbol COL4A1 Species Human, Rat, Avian Reactivity Notes Quail Specificity/Sensitivity Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (BBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NaI of domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1 (beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen alpha1 (IV)-alpha2(IV) triple helix. The epitope is triple helical domain about 900 from the N-terminus Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Isotype	IgG1 Kappa
Buffer SomM Sodium Borate	Conjugate	Janelia Fluor 635
Product Description	Purity	Protein A or G purified
Gene ID 1282	Buffer	50mM Sodium Borate
Gene Symbol COL4A1 Species Human, Rat, Avian Caactivity Notes Quail Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Product Description	
COL4A1	Host	Mouse
Reactivity Notes Quail Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV have been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Gene ID	1282
Reactivity Notes Quail Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen alpha1 (IV)-alpha2(IV) triple helix. The epitope is triple helical domain about 900 from the N-terminus Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Gene Symbol	COL4A1
Specificity/Sensitivity Collagen Type IV is a major component of the basement membrane and plays an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Species	Human, Rat, Avian
an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen. Immunogen Immunogen Immunogen Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Reactivity Notes	Quail
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.		an important role in cell adhesion, migration, differentiation and growth. Collagen Type IV express at the basement membranes in a variety of tissues including kidney, muscle, lymph nodes, lung, tendon and spleen. Collagen Type IV has been shown to be useful in differentiating microinvasive from in situ ductal carcinomas of the breast. Other Collagen Type IV studies include use in pancreatic adenocarcinoma and chronic pancreatitis, nephrosclerosis and other kidney diseases, oral squamous cell carcinoma, laryngeal cancers, ovarian cancers and cervical cancers. Type IV collagen is the major structural component of glomerular basement membranes (GBM), forming a meshwork together with laminins, proteoglycans and entactin/nidogen. Arresten, comprising the C-terminal NC1 domain, inhibits angiogenesis and tumor formation. The C-terminal half is found to possess the anti-angiogenic activity. Specifically inhibits endothelial cell proliferation, migration and tube formation. Inhibits expression of hypoxia-inducible factor 1alpha and ERK1/2 and p38 MAPK activation. Ligand for alpha1/beta1 integrin. M3F7 recognizes type IV collagen in basement membranes in kidney, lung, placenta, cornea and skin. This antibody does not recognize denatured type IV collagen.
Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	Immunogen	
Product Application Details	Notes Product Application Details	

Product Application Details

Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin **Applications**



Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin
Application Notes	Ontimal dilution of this antibody should be experimentally determined



Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

Application Notes

nb-customerservice@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08515JF635

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

