## **Product Datasheet**

### Cytokeratin 14 Antibody (KRT14/2375) [Alexa Fluor® 594] NBP3-08531AF594

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08531AF594

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08531AF594



#### NBP3-08531AF594

Cytokeratin 14 Antibody (KRT14/2375) [Alexa Fluor® 594]

basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Cytokeratin 14 Antibody (KK114/2373) [Alexa Fluor® 394]		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneKRT14/2375Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKRT14HostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneKRT14/2375Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Unit Size	100 ul	
ClonalityMonoclonalCloneKRT14/2375Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is one of the specific is and markers for distinguishing	Concentration		
CloneKRT14/2375Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, some glandular epithelia, some of the specific basal markers for distinguishing	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Clonality	Monoclonal	
IsotypeIgG2a KappaConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Clone	KRT14/2375	
ConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Isotype	IgG2a Kappa	
Buffer 50mM Sodium Borate   Product Description Mouse   Host Mouse   Gene ID 3861   Gene Symbol KRT14   Species Human, Mouse, Rat, Porcine   Marker Squamous Cell Marker   Specificity/Sensitivity Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Conjugate	Alexa Fluor 594	
Product DescriptionHostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Purity	Protein A or G purified	
HostMouseGene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Buffer	50mM Sodium Borate	
Gene ID3861Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Product Description		
Gene SymbolKRT14SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Host	Mouse	
SpeciesHuman, Mouse, Rat, PorcineMarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Gene ID	3861	
MarkerSquamous Cell MarkerSpecificity/SensitivityCytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Gene Symbol	KRT14	
Specificity/Sensitivity Cytokeratin 14 (CK14) belongs to the type I (or A or acidic) subfamily of low molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Species	Human, Mouse, Rat, Porcine	
molecular weight keratins and exists in combination with keratin 5 (type II or B basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing	Marker	Squamous Cell Marker	
as well as for differentiation of benign prostate from prostate carcinoma.	Specificity/Sensitivity	molecular weight keratins and exists in combination with keratin 5 (type II or B or basic). CK14 is found in basal cells of squamous epithelia, some glandular epithelia, myoepithelium, and mesothelial cells. Anti-CK14 is useful in differentiating squamous cell carcinomas from poorly differentiated epithelial tumors. Anti-CK14 is one of the specific basal markers for distinguishing between basal and non-basal subtypes of breast carcinomas. Anti-CK14 is also a good marker for differentiation of intraductal from invasive salivary duct carcinoma by the positive staining of basal cells surrounding the in-situ neoplasm as well as for differentiation of benign prostate from prostate carcinoma. Furthermore, this antibody has been useful in separating oncocytic tumors of the kidney from its renal mimics, and in identifying metaplastic carcinomas of the	
ImmunogenRecombinant human Cytokeratin 14 fragment (around aa351-472) (exact sequence is proprietary) (Uniprot: P02533)	Immunogen		



Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an	
academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.	
Product Application Details	
Flow Cytometry, Immunohistochemistry-Paraffin	
Flow Cytometry, Immunohistochemistry-Paraffin	
Optimal dilution of this antibody should be experimentally determined.	

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-08531AF594

NBP1-84917PEP	Cytokeratin 14 Recombinant Protein Antigen
1129-ER-050	ErbB2/Her2 [Unconjugated]
NBP1-72467-100ug	Recombinant Human Cytokeratin 14 His Protein
NBP1-96981AF594	Mouse IgG2a Kappa Isotype Control (M2AK) [Alexa Fluor® 594]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08531AF594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

