## **Product Datasheet**

### SDHB Antibody (SDHB/2382) [Alexa Fluor® 647] NBP3-08537AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08537AF647

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08537AF647



#### NBP3-08537AF647

SDHB Antibody (SDHB/2382) [Alexa Fluor® 647]

Product	Information
---------	-------------

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSDHB/2382Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBufferSom Sodium BorateProduct DescriptionMouseGene IDG390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits complex and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor supressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around an 465-273) of human SDHB protein (exact sequence is proprietary) (Uniprot: P21912)	Product information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSDHB/2382Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description6390Gene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of phaecochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Unit Size	100 ul
ClonalityMonoclonalCloneSDHB/2382Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description6390Gene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex. a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC) and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Concentration	•
CloneSDHB/2382Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function, Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHA), an iron- sulfur protein (SDHB) and two membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron-sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit protein slead to destabilization of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Clonality	Monoclonal
IsotypeIgG2b KappaConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits souffur protein (SDHB) and two membrane-bound subunits (SDHC) and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Clone	SDHB/2382
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC) and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Isotype	IgG2b Kappa
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC) and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Conjugate	Alexa Fluor 647
Product DescriptionHostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Purity	Protein A or G purified
HostMouseGene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Buffer	50mM Sodium Borate
Gene ID6390Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Product Description	
Gene SymbolSDHBSpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Host	Mouse
SpeciesHumanMarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Gene ID	6390
MarkerPheochromocytoma MarkerSpecificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Gene Symbol	SDHB
Specificity/SensitivitySuccinate dehydrogenase (SDH) is Complex II in the mitochondria, vital for mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron- sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.ImmunogenRecombinant fragment (around aa 165-273) of human SDHB protein (exact	Species	Human
<ul> <li>mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron-sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of phaeochromocytomas, paragangliomas and GIST.</li> <li>Immunogen</li> </ul>	Marker	Pheochromocytoma Marker
	Specificity/Sensitivity	mitochondrial electron transport, as well as Krebs cycle function. Four subunits comprise the SDH protein complex: a flavochrome subunit (SDHA), an iron-sulfur protein (SDHB) and two membrane-bound subunits (SDHC and SDHD) anchored to the inner mitochondrial membrane. The SDH complex functions as a tumor suppressor. Loss of any subunit proteins lead to destabilization of the complex and tumor formation. Antibody to SDHB is helpful in the identification of
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Protein Array
<b>Recommended Dilutions</b>	Western Blot, Protein Array
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-08537AF647

NBP1-43317AF647	Mouse IgG2b Kappa Light Chain Isotype Control (MG2b) [Alexa Fluor® 647]
NBP1-87069PEP	SDHB Recombinant Protein Antigen
NBL1-15766	SDHB Overexpression Lysate
AF482	Ret Antibody [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08537AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

