Product Datasheet

Cytokeratin 17 Antibody (KRT17/4604) [Alexa Fluor® 647] NBP3-08541AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08541AF647

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08541AF647



NBP3-08541AF647

Cytokeratin 17 Antibody (KRT17/4604) [Alexa Fluor® 647]

Storage Store at 4C in the dark. Clonality Monoclonal Clone KRT17/4604 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation	Cytokeratii 17 7 iitibody (NNT 1774004) [ritexa 1 idol@ 047]		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone KRT17/4604 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighth five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinomas considered course. Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone KRT17/4604 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in in LADC. Eight five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinomas immunoreactivity represents	Unit Size	100 ul	
Clone KRT17/4604 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighther five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course. Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Clone KRT17/4604 Preservative 0.05% Sodium Azide IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of differentiation. CK17 is expressed in CK17 ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 ab marks basal cell differentiation. CK17 is expressed in epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 ab marks beast carcinoma immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Storage	Store at 4C in the dark.	
Preservative Isotype IgG1 Kappa Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighther five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Clonality	Monoclonal	
Isotype	Clone	KRT17/4604	
Conjugate Alexa Fluor 647 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighther five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Preservative	0.05% Sodium Azide	
Purity Buffer 50mM Sodium Borate Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells or various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Isotype	IgG1 Kappa	
Buffer 50mM Sodium Borate	Conjugate	Alexa Fluor 647	
Product Description Host Mouse Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eight five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Purity	Protein A or G purified	
Host Gene ID 3872 Gene Symbol KRT17 Species Human Marker Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eight five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Buffer	50mM Sodium Borate	
Gene Symbol KRT17 Species Human Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighther five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Product Description		
Gene Symbol KRT17 Species Human Basal Epithelial Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Host	Mouse	
Marker Basal Epithelial Marker Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eight five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Gene ID	3872	
Marker Specificity/Sensitivity Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Gene Symbol	KRT17	
Cytokeratin 17 (CK17) is normally expressed in the basal cells of complex epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course. Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Species	Human	
epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excell tool to distinguish myoepithelial cells from luminal epithelium of various gland such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents	Marker	Basal Epithelial Marker	
intestinal subtype.	Specificity/Sensitivity	epithelia but not in stratified or simple epithelia. Antibody to CK17 is an excellent tool to distinguish myoepithelial cells from luminal epithelium of various glands such as mammary, sweat and salivary. CK17 is expressed in epithelial cells of various origins, such as bronchial epithelial cells and skin appendages. It may be considered as epithelial stem cell marker because CK17 Ab marks basal cell differentiation. CK17 is expressed in SCLC much higher than in LADC. Eighty-five percent of the triple negative breast carcinomas immunoreact with basal cytokeratins including anti-CK17. Also important is that cases of triple negative breast carcinoma with expression of CK17 show an aggressive clinical course. Additionally, anti-CK17 and anti-MUC1 immunoreactivity represents pancreatobiliary subtype whereas anti-MUC2 and anti-CDX-2 positivity defines	
Immunogen Recombinant full-length human Cytokeratin 17 protein (Uniprot: Q04695)	Immunogen	Recombinant full-length human Cytokeratin 17 protein (Uniprot: Q04695)	



Notes

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08541AF647

IC002R Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 647]

NBP2-51594-0.1mg Recombinant Human Cytokeratin 17 His Protein

1503-SE-010 Furin [Unconjugated]

H00003872-T01 Cytokeratin 17 293T Cell Transient Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08541AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

