Product Datasheet

Mitochondria Antibody (MTC02/2860R) [Alexa Fluor® 647] NBP3-08548AF647

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08548AF647

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08548AF647



NBP3-08548AF647

Mitochondria Antibody (MTC02/2860R) [Alexa Fluor® 647]

Product Information

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneMTC02/2860RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondria in cells or specifically with human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondria in cells or specifically with human cells. It can be used to stain mitochondria in cells or specifically with human cells. It is nenographic model research. It reacts specifically with human cells. It is nenographic model research. It reacts specifically with human cells. It is nenographic model research. It reacts specifically with human cells. It is nenoclonal antibody facilitates the complex differential diagnosis of granular renal cell tumors. Reportedy, this monoclonal antibody facilitates the classification of salivary tumors.ImmunogenSemi-Purified mitochondrial preparation	Froduct mormation	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneMTC02/2860RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in numan cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondria in cells or tissue preparations and can be used as a mitochondria marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in tenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. It monoclonal antibody facilitates the classification of salivary tumors.	Unit Size	100 ul
ClonalityMonoclonalCloneMTC02/2860RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells, in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. It monoclonal antibody racilitates the classification of salivary tumors.	Concentration	•
CloneMTC02/2860RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specificially with human cells in neuronal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in cell or tissue preparations and can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondria in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with numan cells in xenographic model research. It reacts specifically with numan cells in xenographic model research. It reacts specifically with numan cells in xenographic model research. It reacts specificatly differential diagnosis of granular renal cell tumors. Reportedy, this monoclonal antibody facilitates the classification of salivary tumors.	Clonality	Monoclonal
IsotypeIgGConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionForein A managementHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in numa cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells in xenographic model research. It reacts specifically with numan cells in a suseful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Clone	MTC02/2860R
ConjugateAlexa Fluor 647PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Isotype	IgG
Buffer50mM Sodium BorateProduct DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody facilitates the classification of salivary tumors.	Conjugate	Alexa Fluor 647
Product DescriptionHostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Purity	Protein A or G purified
HostRabbitSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Buffer	50mM Sodium Borate
SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Product Description	
Reactivity NotesDoes not react with Mouse or Rat.MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Host	Rabbit
MarkerMarker for Human Cells, Granular RCC's & Salivary TumorsSpecificity/SensitivityThis monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Species	Human, Mouse (Negative), Rat (Negative)
Specificity/Sensitivity This monoclonal antibody recognizes a 60kDa antigen associated with the mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Reactivity Notes	Does not react with Mouse or Rat.
mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the classification of salivary tumors.	Marker	Marker for Human Cells, Granular RCC's & Salivary Tumors
Immunogen Semi-Purified mitochondrial preparation	Specificity/Sensitivity	mitochondria in human cells. It can be used to stain mitochondria in cell or tissue preparations and can be used as a mitochondrial marker in subcellular fractions. It produces a spaghetti-like pattern in normal and malignant cells. This antibody is an excellent marker for human cells in xenographic model research. It reacts specifically with human cells, including neurons and embryonic stem cells. Immunostaining pattern with anti-mitochondrial monoclonal antibody has been reported as a useful discriminatory adjunct in the complex differential diagnosis of granular renal cell tumors. Reportedly, this monoclonal antibody facilitates the
	Immunogen	Semi-Purified mitochondrial preparation

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08548AF647

NBP2-24891AF647Rabbit IgG Isotype Control [Alexa Fluor® 647]NBP2-29448Mitochondria Isolation Kit

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08548AF647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

