Product Datasheet

Thyroid Peroxidase Antibody (TPO/3813R) [Alexa Fluor® 594] NBP3-08761AF594

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08761AF594

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08761AF594

NBP3-08761AF594

Thyroid Peroxidase Antibody (TPO/3813R) [Alexa Fluor® 594]

Product InformationUnit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneTPO/3813RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionTPOHostRabbitGene SymbolTPOSpeciesHumanMarkerThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroig dlyand phenoxy-ester formation in the idination of tyrosine residues of thyroid hormones, thyroxine and thyroid hormones, thyroxine and thyroid hormones, thyroxine exhibit an anomaly in TPO resulting on defent the dividio domines of thyroid hormones, thyroxine exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between berign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804) (Exact sequence is proprietary) (Uniprot: PO7202)			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneTPO/3813RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description7173HostRabbitGene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroid MarkerSpecificity/SensitivityThyroid MarkerSpecificity/Sensitivityaley a role in thyroid gland function and autoimmunity. It is present as a dime on the apical surface of thyroid follicular cells. TPO functions in the oid nation of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and thiod thyronine. Mutations in this gene as associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and maligna	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneTPO/3813RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionTPOHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpeciesThyroid MarkerSpecificity/SensitivityThyroid set of thyroid follcular cells. TPO functions in the iodination of tyrosine to gene as associated with several disorders of thyroid hormones, thyroxine and tridodthyronine. Mutations in this gene are associated with several disorders of thyroid hormone presention of the origine set one present on the apical surface of thyroid hormones, thyroxine and tridodthyronine. Mutations in this gene are associated with several disorders of thyroid hormone presis. Including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between being and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between being and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between being and malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between being and malignant thyroid tumor	Unit Size	100 ul	
ClonalityMonoclonalCloneTPO/3813RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpeciesHumanSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO furctions in the loidination of tyrosine is in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and tridothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors. exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between being and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Concentration		
CloneTPC/3813RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBufferSomM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid folloular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid more, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between beingn and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Clonality	Monoclonal	
IsotypeIgGConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triidothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors. Recombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Clone	TPO/3813R	
ConjugateAlexa Fluor 594PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Isotype	IgG	
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Conjugate	Alexa Fluor 594	
Product DescriptionHostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Purity	Protein A or G purified	
HostRabbitGene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Buffer	50mM Sodium Borate	
Gene ID7173Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonegenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Product Description		
Gene SymbolTPOSpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Host	Rabbit	
SpeciesHumanMarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Gene ID	7173	
MarkerThyroid MarkerSpecificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Gene Symbol	TPO	
Specificity/SensitivityThyroperoxidase (TPO) is a 933-amino acid, type I transmembrane glycoprotein that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Species	Human	
that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody may aid in the differentiation between benign and malignant thyroid tumors.ImmunogenRecombinant fragment of human Thyroid Peroxidase (around aa 685-804)	Marker	Thyroid Marker	
	Specificity/Sensitivity	that plays a key role in thyroid gland function and autoimmunity. It is present as a dimer on the apical surface of thyroid follicular cells. TPO functions in the iodination of tyrosine residues in thyroglobulin and phenoxy-ester formation between pairs of iodinated tyrosines to generate the thyroid hormones, thyroxine and triiodothyronine. Mutations in this gene are associated with several disorders of thyroid hormonogenesis, including congenital hypothyroidism, congenital goiter, and thyroid hormone organification defect IIA. Malignant thyroid tumors exhibit an anomaly in TPO resulting in lower affinity for anti-TPO. This antibody	
	Immunogen		

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08761AF594

IC1051T	Rabbit IgG Isotype Control (60024B) [Alexa Fluor® 594]
H00007173-P01-2ug	Recombinant Human Thyroid Peroxidase GST (N-Term) Protein
210-TA-005	TNF-alpha [Unconjugated]
H00007173-Q01-10ug	Recombinant Human Thyroid Peroxidase GST (N-Term) Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08761AF594

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

