Product Datasheet

RBM8A Antibody (PCRP-RBM8A-1B4) [Alexa Fluor® 488] NBP3-08800AF488

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08800AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08800AF488



NBP3-08800AF488

RBM8A Antibody (PCRP-RBM8A-1B4) [Alexa Fluor® 488]

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalClonePCRP-RBM8A-1B4Preservative0.05% Sodium AzideIsotypeIgG1ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID9939Gene SymbolRBM8ASpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated raman and Xenopus laevis origin. RBM8A Antibody is a hultipe origing agreement of exon- pusch and Xenopus laevis origin. RBM8A Antibody form, as well as multiple complex that assembles approximately 20-24 nucleotides upstream of exon- complex that assembles approximately 20-24 nucleotides upstream of exon- scon junctions in pre-mRNs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC, include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, Aly/REF, Magoh, ANREF, and Magoh have the ability bo communicate to the cytoplasm the processing history of the mRNA, including the position of the removed intons. The gene encoding human RBM8A encodes three transcripts. RBM8A A is a ubiquitously expres	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone PCRP-RBM8A-1B4 Preservative 0.05% Sodium Azide Isotype IgG1 Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description House Gene ID 9939 Gene Symbol RBM8A Species Hurman Specificity/Sensitivity RBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated roms of anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody 10. including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex flat assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nosense-mediated mRNA decay. Members of the EJC include RBM8A, Alv/REF, Magoh, RNPS1, SRm160, and DEK Alv/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNA sin the cytoplasm until they are translated. Therefore, RBM8A, Alv/REF, Magoh, and RBM8A, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously servessed protein. Although RBM8A is complex that the costore of the cell.	Unit Size	100 ul
ClonalityMonoclonalClonePCRP-RBM8A-1B4Preservative0.05% Sodium AzideIsotypeIgG1ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody form, as well as multiple conjugated form, and Nanopus Levis origin. RBM8A Antibody (also designated RBM8A antibody), including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A, antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junction, and nonsnes-mediated mRNA exopt, cytoplasmic localization, and nonsnes-mediated mRNA exopt, mothes in the cytoplasmic localization, and nonsnes-mediated mRNA exopt ments, and these proteins remain stably bound to spliced mRNA is in the cytoplasmic localization, and nonsnes-mediated mRNA exopt, methys in the cytoplasmic localization is the cytoplasm. The gene encoding human RBM8A, nicluding the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNA, including the position of the cremoved introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is	Concentration	•
ClonePCRP-RBM8A-1B4Preservative0.05% Sodium AzideIsotypeIgG1ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated formouse, rat, human and Xenopus laevis origin. RBM8A antibody (ac) is a valiable as both the non-conjugated anti-RBM8A antibody including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (LJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the ELO, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, Magoh, and RBM8A, Aly/REF, Magoh, and SBM8A, Aly/REF, Magoh, Aly/REF, Magoh, and SBM8A, Aly/REF, Magoh, Magoh, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm. The predominantly detected in the nucleus and is co-localized with oskar mRNA at the posterior pole of the cell.	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG1ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monochonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A antibody (form, as well as multiple conjugated forms of anti-RBM8A antibody (form, as well as multiple conjugated forms of anti-RBM8A antibody (consplex (LC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exonexon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, Ald RBM8A, Aly/REF, Magoh, Ald RBM8A, Aly/REF, Magoh, Ald RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK, Aly/REF, Magoh, and RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK, Aly/REF, Magoh, and hese proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.	Clonality	Monoclonal
IsotypeIgG1ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm util they are transcripts. RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8A is a ubiquitously expressed protein. Although RBM8A, shuttles to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitouyle yaressed protein. Although RBM8A, shuttles to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitouyle yaressed protein. Although RBM8A shuttles to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitouyle yaressed protein. Although RBM8A shuttles to the cytoplasm the processing h	Clone	PCRP-RBM8A-1B4
ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotide supstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, Magoh, And RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and these proteins remain stably bound to spliced mRNAs in the cytoplasm communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody, form, as well as multiple conjugated forms of anti-RBM8A antibody, involuting agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Isotype	IgG1
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody form, as well as multiple conjugated forms of anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody 20-24 nucleotides upstream of exonexon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC, and RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.	Conjugate	Alexa Fluor 488
Product DescriptionHostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm, the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.	Purity	Protein A or G purified
HostMouseGene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A Antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody (also designated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, digentified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Buffer	50mM Sodium Borate
Gene ID9939Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Product Description	
Gene SymbolRBM8ASpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK, Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Host	Mouse
SpeciesHumanSpecificity/SensitivityRBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Gene ID	9939
Specificity/Sensitivity RBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.	Gene Symbol	RBM8A
RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon- exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co- localized with oskar mRNA at the posterior pole of the cell.	Species	Human
Immunogen Recombinant full-length human RBM8A protein (Uniprot: Q9Y5S9)	Specificity/Sensitivity	RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.
	Immunogen	Recombinant full-length human RBM8A protein (Uniprot: Q9Y5S9)



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Protein Array
Recommended Dilutions	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Protein Array
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08800AF488

DDXCM01A488	Mouse IgG1 Isotype Control [Alexa Fluor® 488]
NBP1-51060-0.05mg	Recombinant Human RBM8A His Protein
DPSG10	PSG1 [HRP]
NBL1-15217	RBM8A Overexpression Lysate

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08800AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

