Product Datasheet

RBM8A Antibody (PCRP-RBM8A-1B4) [mFluor Violet 610 SE] NBP3-08800MFV610

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08800MFV610

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-08800MFV610



NBP3-08800MFV610

RBM8A Antibody (PCRP-RBM8A-1B4) [mFluor Violet 610 SE]

0.1 ml
0.1 ml
Please see the vial label for concentration. If unlisted please contact technical services.
Store at 4C in the dark.
Monoclonal
PCRP-RBM8A-1B4
0.05% Sodium Azide
IgG1
mFluor Violet 610 SE
Protein A or G purified
50mM Sodium Borate
Mouse
9939
RBM8A
Human
RBM8A Antibody is a high quality monoclonal RBM8A antibody (also designated RBM8A antibody) suitable for the detection of the RBM8A protein of mouse, rat, human and Xenopus laevis origin. RBM8A Antibody (4C4) is available as both the non-conjugated anti-RBM8A antibody form, as well as multiple conjugated forms of anti-RBM8A antibody, including agarose, HRP, PE, FITC and multiple Alexa Fluor conjugates. The exon junction complex (EJC) is a multiprotein complex that assembles approximately 20-24 nucleotides upstream of exon-
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is co-
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell.
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell. Recombinant full-length human RBM8A protein (Uniprot: Q9Y5S9) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell. Recombinant full-length human RBM8A protein (Uniprot: Q9Y5S9) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The
exon junctions in pre-mRNAs. It is involved in mRNA export, cytoplasmic localization, and nonsense-mediated mRNA decay. Members of the EJC include RBM8A, Aly/REF, Magoh, RNPS1, SRm160, and DEK. Aly/REF, Magoh, and RBM8A, identified as RBM8 in mouse and rat, make up the core of the EJC, and these proteins remain stably bound to spliced mRNAs in the cytoplasm until they are translated. Therefore, RBM8A, Aly/REF, and Magoh have the ability to communicate to the cytoplasm the processing history of the mRNA, including the position of the removed introns. The gene encoding human RBM8A encodes three transcripts. RBM8A is a ubiquitously expressed protein. Although RBM8A shuttles to the cytoplasm, it is predominantly detected in the nucleus and is colocalized with oskar mRNA at the posterior pole of the cell. Recombinant full-length human RBM8A protein (Uniprot: Q9Y5S9) mFluor(TM) is a trademark of AAT Bioquest, Inc. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence,



Images

RBM8A Antibody (PCRP-RBM8A-1B4) [mFluor Violet 610 SE] [NBP3-08800MFV610] - Vial of mFluor Violet 610 conjugated antibody. mFluor Violet 610 is optimally excited at 421 nm by the Violet laser (405 nm) and has an emission maximum of 613 nm.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-08800MFV610

NBP1-51060-0.05mg Recombinant Human RBM8A His Protein

DPSG10 PSG1 [HRP]

NBL1-15217 RBM8A Overexpression Lysate

H00023435-M01 TDP-43/TARDBP Antibody (2E2-D3)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08800MFV610

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

