Product Datasheet

Bcl-6 Antibody (BCL6/1718) [FITC] NBP3-08865F

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08865F

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08865F



NBP3-08865F

Bcl-6 Antibody (BCL6/1718) [FITC]

| Product Information | |
|-----------------------------|--|
| Unit Size | 100 ul |
| Concentration | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage | Store at 4C in the dark. |
| Clonality | Monoclonal |
| Clone | BCL6/1718 |
| Preservative | 0.05% Sodium Azide |
| Isotype | IgG1 Kappa |
| Conjugate | FITC |
| Purity | Protein A or G purified |
| Buffer | PBS |
| Product Description | |
| Host | Mouse |
| Gene ID | 604 |
| Gene Symbol | BCL6 |
| Species | Human |
| Marker | Follicular Lymphoma Marker |
| Specificity/Sensitivity | Recognizes a protein of 95kDa, which is identified as Bcl-6. Antibody to bcl-6 is helpful in a number of diagnostic settings: (1) In the differential diagnosis of small B-cell lymphoma. Follicular lymphoma will show bcl-6 (and CD10) positivity whereas other small B-cell lymphomas are usually negative. (2) Bcl-6 is an important prognostic marker in diffuse large B-cell lymphomas (DLBCL), where CD10, bcl-6 and MUM1/IRF4 are used to identify germinal center and activated B-cell phenotypes. (3) Bcl-6 can be valuable in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL). The Reed-Sternberg cells of classical Hodgkin lymphoma are bcl-6 negative whereas the large ('LH') cells of NLPHL are bcl-6 positive. In contrast, anti-Bcl-6 rarely stains mantle-cell lymphoma and MALT lymphoma. |
| Immunogen | Recombinant human bcl-6 protein fragment (aa256-389) (exact sequence is proprietary) (Uniprot: P41182) |
| Product Application Details | |
| Applications | Immunohistochemistry-Paraffin, Protein Array |
| Recommended Dilutions | Immunohistochemistry-Paraffin, Protein Array |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08865F

| NBP1-43319F | Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [FITC] |
|--------------------|--|
| H00000604-Q01-10ug | Recombinant Human Bcl-6 GST (N-Term) Protein |
| 594-ML-010 | IL-21 [Unconjugated] |
| NBL1-07951 | Bcl-6 Overexpression Lysate |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08865F

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

