Product Datasheet

Von Willebrand Factor Antibody (rVWF/2480) [Janelia Fluor® 646] NBP3-08944JF646

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08944JF646

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08944JF646

NBP3-08944JF646

Von Willebrand Factor Antibody (rVWF/2480) [Janelia Fluor® 646] Product Information Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone rVWF/2480 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer SomM Sodium Borate Product Description Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial and modifications that influence the affinity and availability for Factor VIII and promotes platelet achesion and aggregation. WWF undergoes a variety of UII, including cleavage of the propeptide and formation of N-terminal disulfide body helps to establish the endothelial availability for Factor VIII, including cleavage of othe propeptide and formation of N-terminal disulfide body helps to establish the endothelial antiferentiation or within a panel of other vascular markers although not alt lumors of endothelial differentiating vascular lesions from those of other tissue			
Unit Size 100 ul Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clona rVWF/2480 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Janelia Fluor 646 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Wilebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. VWF undergoes a variety of postranslational modifications that influence the affinity and availability for Factor VIII including cleavage of the propertie and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial anature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiation vacular markers atthough not all tumors of endothelial flature of endothelial flature of some lesions of	Von Willebrand Factor Antibody (rVWF/2480) [Janelia Fluor® 646]		
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloner/WF/2480Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. WF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. (Kaposi S sarcom and cardiac myxoma, It is widely used for differentiating vascular lesions for those of other tissue differentiation within a panel of other vascular markers athough not all tumors of endothelial differentiation contain this sartigon.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalClonerVWF/2480Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description7450HostMouseGene ID7450SpeciesHumanMarkerEndothelial MarkerSpeciesVWFSpeciesvon Willebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of vised for differentiating vascular lesions from those of other tissue differentiation owthin a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes	Unit Size	100 ul	
ClonalityMonoclonalClonerVWF/2480Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. WF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial alture of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation sold under license from the Howard Hughes Medical Institute, Janelia ResearchNotesSold under license from the Howard Hughes Medical Institute, Janelia Research	Concentration	•	
ClonerVWF/2480Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (WF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. WF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial alture of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation within a panel of other vascular lesions from those of other tissue differentiation differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor vIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) NotesNotesSold under license from the Howard Hughes Medical Institute, Janelia Research	Clonality	Monoclonal	
IsotypeIgG1 KappaConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal dissulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s parcoma and cardiac myxoma. It is widely used for differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Clone	rVWF/2480	
ConjugateJanelia Fluor 646PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of postranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to estabilish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular markers although not all tumors of endothelial differentiation contain this antigen. Immunogen Recombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	Isotype	IgG1 Kappa	
Buffer 50mM Sodium Borate Product Description Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial neture of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen. Immunogen Recombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	Conjugate	Janelia Fluor 646	
Product Description Host Mouse Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen. Immunogen Recombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	Purity	Protein A or G purified	
HostMouseGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Buffer	50mM Sodium Borate	
Gene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Product Description		
Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Host	Mouse	
SpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Gene ID	7450	
MarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Gene Symbol	VWF	
Specificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.ImmunogenRecombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Sold under license from the Howard Hughes Medical Institute, Janelia Research	Species	Human	
 endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen. Immunogen Recombinant full-length human Von Willebrand Factor protein (Uniprot: P04275) Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research 	Marker	Endothelial Marker	
Notes Sold under license from the Howard Hughes Medical Institute, Janelia Research	Specificity/Sensitivity	endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi s sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial	
Sold under license from the Howard Hughes Medical Institute, Janelia Research	Immunogen	Recombinant full-length human Von Willebrand Factor protein (Uniprot: P04275)	
	Notes		

Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08944JF646

NBP2-34494PEP	Von Willebrand Factor Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
KA0512	Human Von Willebrand Factor ELISA Kit (Colorimetric)
D6050	IL-6 [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08944JF646

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

