

Product Datasheet

p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free NBP3-08961

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-08961

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-08961



NBP3-08961

p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free


Product Information	
Unit Size	100 ug
Concentration	1 mg/ml
Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	CIP1/4377R
Preservative	No Preservative
Isotype	IgG
Purity	Protein A or G purified
Buffer	10 mM PBS
Target Molecular Weight	21 kDa

Product Description	
Description	1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA & azide. Also available at 200 ug/ml WITH BSA & azide (NBP3-07215). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.
Host	Rabbit
Gene ID	1026
Gene Symbol	CDKN1A
Species	Human
Specificity/Sensitivity	This monoclonal antibody recognizes a 21kDa protein, identified as the p21WAF1 tumor suppressor protein. This monoclonal antibody is highly specific to p21 and shows no cross-reaction with other closely related mitotic inhibitors. p21WAF1 is a specific inhibitor of cdk s and a tumor suppressor involved in the pathogenesis of a variety of malignancies. The expression of this gene acts as an inhibitor of the cell cycle during G1 phase and is tightly controlled by the tumor suppressor protein p53. Its expression is induced by the wild type, but not mutant, p53 suppressor protein. Normal cells generally display a rather intense nuclear p21 expression. Loss of p21 expression has been reported in many carcinomas (gastric carcinoma, non-small cell lung carcinoma, thyroid carcinoma).
Immunogen	Synthetic peptide corresponding to p21/CIP1/CDKN1A residues within aa1-100 of p21/CIP1/CDKN1A was used as an immunogen. (Uniprot: P38936)


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Protein Array
Recommended Dilutions	Western Blot, Immunohistochemistry-Paraffin 1-2 ug/ml, Protein Array
Application Notes	Immunohistochemistry Formalin-fixed: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images


Western Blot: p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free [NBP3-08961] - Western Blot Analysis of HeLa cell lysate using p21/CIP1/CDKN1A antibody (CIP1/4377R).




Immunohistochemistry-Paraffin: p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free [NBP3-08961] - Formalin fixed paraffin embedded human lung carcinoma stained with p21/CIP1/CDKN1A Rabbit Recombinant Monoclonal Antibody (CIP1/4377R).




Immunohistochemistry-Paraffin: p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free [NBP3-08961] - Formalin fixed paraffin embedded human colon carcinoma stained with p21/CIP1/CDKN1A Rabbit Recombinant Monoclonal Antibody (CIP1/4377R).



Immunohistochemistry-Paraffin: p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free [NBP3-08961] - IHC analysis of formalin-fixed, paraffin-embedded human brain. Negative tissue control using p21/CIP1/CDKN1A antibody (CIP1/4377R) at 2ug/ml in PBS for 30min RT. HIER: Tris/EDTA, pH9.0, 45min. 2 : HRP-polymer, 30min. DAB, 5min.



Protein Array: p21/CIP1/CDKN1A Antibody (CIP1/4377R) - Azide and BSA Free [NBP3-08961] - Analysis of Protein Array containing more than 19,000 full-length human proteins using p21/CIP1/CDKN1A Antibody (CIP1/4377R).





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-08961

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-22976	Recombinant Human p21/CIP1/CDKN1A His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08961

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

