Product Datasheet

Von Willebrand Factor Antibody (VWF/1859R) [HRP] NBP3-08963H

Unit Size: 100 ul

Store at 4C in the dark.

www.novusbio.com

G

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-08963H

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-08963H



NBP3-08963H

Von Willebrand Factor Antibody (VWF/1859R) [HRP]

Unit Size100 ulConcentrationPlease see the vial label for concentration. If unlisted please contact tech services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneVWF/1859RPreservativeNo PreservativeIsotypeIgGConjugateHRPPurityProtein A purifiedBufferPBSProduct DescriptionTest RabbitGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in or of disputed histogenesis, e.g. Kaposis sarcom and cardiac myxoma. It is word disputed histogenesis, e.g. Kaposis sarcom and cardiac myxoma. It is word disputed histogenesis, e.g. Kaposis sarcom and cardiac myxoma. It is word disputed histogenesis, e.g. Kaposis sarcom and cardiac myxoma. It is word disputed histogenesis, e.g. Kaposis and cardiac myxoma. It is word disputed histogenesis, e.g. Kaposis and cardiac myxoma. It is word disputed relative the the interimetion of other tissue differentiation of the propertiem at the other listing and cardiac myxoma. It is word the marker satthough not all tumors of endoth			
Storage Store at 4C in the dark. Clonality Monoclonal Clone VWF/1859R Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein A purified Buffer PBS Product Description Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability fo VIII, including cleavage of the propeptide and formation of N-terminal diss bonds. This antibody helps to establish the endothelial nature of some les disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is wuised for differentiating vascular markers although not all tumors of endott			
Clonality Monoclonal Clone VWF/1859R Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein A purified Buffer PBS Product Description Home Host Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal diss bonds. This antibody helps to establish the endothelial nature of some lest disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is we used for differentiating vascular lesions from those of other tissue differenting vascular lesi	nical		
Clone VWF/1859R Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein A purified Buffer PBS Product Description Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some leas disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue differer within a panel of other vascular markers although not all tumors of endoth			
Preservative No Preservative Isotype IgG Conjugate HRP Purity Protein A purified Buffer PBS Product Description Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some les disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all turnors of endotf			
IsotypeIgGConjugateHRPPurityProtein A purifiedBufferPBSProduct DescriptionHostHostRabbitGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some lees disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is wurde of other vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endoth			
ConjugateHRPPurityProtein A purifiedBufferPBSProduct DescriptionRabbitHostRabbitGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac mxyxoma. It is wu used for differentiating vascular lesions from those of other tissue differentiating vascular lesions from those of other tissue differentiating vascular markers although not all tumors of endothered to the propertide and promotes platelet and promotes platelet bits and promotes platelet bits and promotes platelet bits of the translational modifications that influence the affinity and availability for VIII, including cleavage of the propertide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is wu used for differentiating vascular lesions from those of other tissue differentiating vascular lesions from those of other tissue differentiating vascular lesions from those of other tissue differentiating vascular markers although not all tumors of endotheliat tumors of endothe			
Purity Protein A purified Buffer PBS Product Description Rabbit Host Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is wused for differentiating vascular lesions from those of other tissue differer within a panel of other vascular markers although not all tumors of endott			
Buffer PBS Product Description Rabbit Host Rabbit Gene ID 7450 Gene Symbol VWF Species Human Marker Endothelial Marker Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some lest disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue differentiating vascular lesions from those of other tissue differentiating vascular markers although not all tumors of endotter			
Product DescriptionHostRabbitGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some lest disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endott			
HostRabbitGene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue differentiating vascular lesions from those of other tissue differentiating vascular markers although not all tumors of endotted			
Gene ID7450Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varier posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal dist bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endoth	Product Description		
Gene SymbolVWFSpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some lest disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endother			
SpeciesHumanMarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endother			
MarkerEndothelial MarkerSpecificity/Sensitivityvon Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varier posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endother			
Specificity/Sensitivity von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varied posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disubonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is wused for differentiating vascular lesions from those of other tissue differentiating within a panel of other vascular markers although not all tumors of endothered.			
endothelial cells, plasma and platelets. It acts as a carrier protein for Fact and promotes platelet adhesion and aggregation. vWF undergoes a varie posttranslational modifications that influence the affinity and availability for VIII, including cleavage of the propeptide and formation of N-terminal disc bonds. This antibody helps to establish the endothelial nature of some less disputed histogenesis, e.g. Kaposis sarcoma and cardiac myxoma. It is w used for differentiating vascular lesions from those of other tissue different within a panel of other vascular markers although not all tumors of endother			
differentiation contain this antigen.	ty of r Factor ilfide ions of idely tiation		
ImmunogenRecombinant fragment of human Von Willebrand Factor protein (aa1815- (exact sequence is proprietary) (Uniprot: P04275)	1939)		
Product Application Details			
Applications Western Blot, Immunohistochemistry-Paraffin			
Recommended Dilutions Western Blot, Immunohistochemistry-Paraffin			
Application NotesOptimal dilution of this antibody should be experimentally determined.			





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-08963H

NBP2-24891H	Rabbit IgG Isotype Control [HRP]
NBP1-84761PEP	Von Willebrand Factor Recombinant Protein Antigen
210-TA-005	TNF-alpha [Unconjugated]
KA0512	Human Von Willebrand Factor ELISA Kit (Colorimetric)

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-08963H

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

