

# Product Datasheet

## CD96 Antibody (6A6) NBP3-11254

Unit Size: 50 ug

Store at 4C.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP3-11254](http://www.novusbio.com/NBP3-11254)

Updated 1/15/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP3-11254](http://www.novusbio.com/reviews/destination/NBP3-11254)



**NBP3-11254**

CD96 Antibody (6A6)

| Product Information |   |
|---------------------|---|
| Unit Size           | 50 ug                                     |
| Concentration       | 0.1 mg/ml                                 |
| Storage             | Store at 4C.                              |
| Clonality           | Monoclonal                                |
| Clone               | 6A6                                       |
| Preservative        | No Preservative                           |
| Isotype             | IgG2a                                     |
| Purity              | Protein G purified                        |
| Buffer              | 0.2 um filtered solution in PBS, 0.1% BSA |

| Product Description |  |
|---------------------|--|
| Host                | Rat  |
| Gene ID             | 10225  |
| Gene Symbol         | CD96   |
| Species             | Mouse  |
| Immunogen           | The monoclonal antibody 6A6 was raised after immunization of rats with purified recombinant mCD96-hlgG1. |
| Endotoxin Note      | <24 EU/mg  |

| Product Application Details |   |
|-----------------------------|---|
| Applications                | Western Blot, Flow Cytometry, Functional, Immunoassay   |
| Recommended Dilutions       | Western Blot, Flow Cytometry, Functional, Immunoassay   |
| Application Notes           | For Western blotting and Flow Cytometry dilutions to be used depend on detection system applied. It is recommended that users test the reagent and determine their own optimal dilutions. The typical starting working dilution is 1:50. For functional studies dilutions have to be made according to the amounts of CD96 to be inactivated. |





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-  
techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP3-11254](http://www.novusbio.com/reviews/submit/NBP3-11254)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

