## **Product Datasheet**

### MUC5AC Antibody (2-11M1) [Janelia Fluor® 525] NBP3-11418JF525

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11418JF525

Updated 8/20/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11418JF525



#### NBP3-11418JF525

MUC5AC Antibody (2-11M1) [Janelia Fluor® 525]

| , , , , , , , , , , , , , , , , , , , |  |
|---------------------------------------|--|
| Product Information                   |  |
| Unit Size                             | 0.1 ml   |
| Concentration                         | Please see the vial label for concentration. If unlisted please contact technical services.  |
| Storage                               | Store at 4C in the dark.   |
| Clonality                             | Monoclonal   |
| Clone                                 | 2-11M1   |
| Preservative                          | 0.05% Sodium Azide   |
| Isotype                               | IgG1 Kappa   |
| Conjugate                             | Janelia Fluor 525  |
| Purity                                | Protein A or G purified  |
| Buffer                                | 50mM Sodium Borate   |
| Product Description                   |  |
| Host                                  | Mouse  |
| Gene ID                               | 4586   |
| Gene Symbol                           | MUC5AC   |
| Species                               | Human, Mouse, Bovine, Feline, Monkey   |
| Specificity/Sensitivity               | This monoclonal antibody recognizes the peptide core of gastric mucin M1 (recently identified as Mucin 5AC).Its epitope is located in the N-terminal cysteine rich part of the peptide core of MUC5AC, which is heavily glycosylated. Its epitope is destroyed by beta-mercaptoethanol but not by periodate treatment. monoclonal antibody 2-11M1 reacts with the protein backbone exclusively; it only reacts with fully deglycosylated MUC5AC. Therefore, the material under test should also be fully deglycosylated. This can be achieved with standard periodate oxidation method. The success of the deglycosylation can be checked with routine PAS (Periodic Acid Shiff) staining. After deglycosylation, the preparation should no longer be stainable with PAS reagent. Only then 2-11M1 will react should MUC5AC be present. This mucin is present in primary ovarian mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies, Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well as in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas. |
| Immunogen                             | M1 mucin preparation from the fluid of an ovarian mucinous cyst belonging to an O Le(a-b) patient  |
| Notes                                 | Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.  |
| Product Application Details           |  |
| Applications                          | ELISA, Flow Cytometry, Immunocytochemistry/ Immunofluorescence,<br>Immunofluorescence  |
| Recommended Dilutions                 | Flow Cytometry, ELISA, Immunocytochemistry/ Immunofluorescence,<br>Immunofluorescence  |
|                                       |  |



| Application Notes | Optimal dilution of this antibody should be experimentally determined. |
|-------------------|--|
|-------------------|--|





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-11418JF525

| H00004586-Q01-10ug | Recombinant Human MUC5AC GST (N-Term) Protein |
|--------------------|---|
| 210-TA-005         | TNF-alpha [Unconjugated]                      |
| NBP2-76703         | Human MUC5AC ELISA Kit (Colorimetric)         |
| 236-EG-200         | EGF [Unconjugated]                            |

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11418JF525

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

