# **Product Datasheet**

# Macrophage and Histiocytoma Marker Antibody (D11) [Janelia Fluor® 669] NBP3-11429JF669

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11429JF669

Updated 8/20/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-11429JF669



## NBP3-11429JF669

Macrophage and Histiocytoma Marker Antibody (D11) [Janelia Fluor® 669]		
Product Information		
Unit Size	0.1 ml	
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.	
Storage	Store at 4C in the dark.	
Clonality	Monoclonal	
Clone	D11	
Preservative	0.05% Sodium Azide	
Isotype	IgG1 Kappa	
Conjugate	Janelia Fluor 669	
Purity	Protein A or G purified	
Buffer	50mM Sodium Borate	
Product Description		
Host	Mouse	
Species	Human, Mouse (Negative), Porcine (Negative), Rat (Negative)	
Reactivity Notes	Does not react with Mouse, Rat or Porcine.	
Marker	Macrophage & Histiocytoma Marker	
Specificity/Sensitivity	In Western blotting, it detects an antigen of 125kDa in human liver and 135kDa in tumors of histiocytic origin. Comparative study of this monoclonal antibody and a standard CD68 monoclonal antibody showed that their antigens are different. Its antigen in all macrophage types studied is located on the plasma membrane and within cytoplasmic structures including lysosomes. This monoclonal antibody shows a restricted reactivity to cells of the monocyte/macrophage system. It specifically reacts with blood monocytes and stains resident macrophages in a wide variety of human tissues. This monoclonal antibody does not stain antigenpresenting cells, e.g., Langerhans cells. Reportedly, its reactivity is restricted to histiocytes and macrophages.	
Immunogen	Membrane preparation from human hepatocytes	
Notes	Sold under license from the Howard Hughes Medical Institute, Janelia Research Campus.	
Product Application Details		
Applications	Immunohistochemistry, Immunohistochemistry-Paraffin	
Recommended Dilutions	Immunohistochemistry, Immunohistochemistry-Paraffin	

Necommended Dilutions	mindronistochemistry, mindronistochemistry-i aramir
Application Notes	Optimal dilution of this antibody should be experimentally determined.





### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11429JF669

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

