Product Datasheet

EpCAM/TROP1 Antibody (SPM491) [CoraFluor™ 1] NBP3-11478CL1

Unit Size: 0.1 ml

Store at 4C in the dark. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11478CL1

Updated 10/22/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11478CL1



NBP3-11478CL1

EpCAM/TROP1 Antibody (SPM491) [CoraFluor™ 1]

Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays.HostMouseGene ID4072Gene SymbolEPCAMSpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenRecombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422)		
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Do not freeze. Clonality Monoclonal Clone SPM491 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 430 nm, and emits at approximately 490 nm, and be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial specf	Product Information	
services. Storage Store at 4C in the dark. Do not freeze. Cionality Monoclonal Cione SPM491 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(TM) 1 absorbs UV light at approximately 40 orm. and emits at approximately 40 orm. ande approximately 40 orm. and emits at approximately	Unit Size	0.1 ml
Clonality Monoclonal Clone SPM491 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emission ma 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 caneFluor(TM) 1 caneFluor(IM) 1 absorbs UV light at approximately 340 nm, and emission acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 caneFluor(TM) 1 coneFluor(TM) 1 caneFluor(TM) 1	Concentration	•
Cione SPM491 Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolv Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence Resonance Energy Transfer) Host Mouse Gene ID 4072 Gene Symbol EPCAM Specificity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensit	Storage	Store at 4C in the dark. Do not freeze.
Preservative No Preservative Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 Description CoraFluor(IM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 Soborb XI: bight at approximately 340 nm and emits at approximately 440 nm 545 nm ,585 nm and 620 nm. It is compatible with common acceptor dyes that aborb at the emission avaelengths of CoraFluor(TM) 1. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial opluar adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adencoarcinoma from pleural mesothelioma and hepatocollular carcinomas. This antibody has been use	Clonality	Monoclonal
Isotype IgG1 Kappa Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 aborbs UV light at approximately 340 nm, and emits at approximately 490 nm s44 sn m, 545 nm, 365 nm and 620 nm. It is compatible with common acceptor dyes that aborb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular achesion molecule (Ep-CAM), it is expressed on baso-lateral cell surface in most simple epithelia and antibody is also userlui in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422) Notes CoraFluor (Clone	SPM491
Conjugate CoraFluor 1 Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) econome Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm s45 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CaraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 p	Preservative	No Preservative
Purity Protein A or G purified Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) door for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 2. Souten 2. CoraFluor(TM) 2. Souten 2. CoraFluor(TM) 2. S	Isotype	IgG1 Kappa
Buffer PBS Product Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(IM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Marker CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Ap	Conjugate	CoraFluor 1
Product Description Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 that absorb of the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody has been used to distinguish as adenocarcinoma from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Applications Western Blot, Flow Cytometry, Immunobistochemistry/Immunofluorescence, and use at aperoscience, and the patentific, Immunofluorescence, anotheory has beneratifin, Immunofluorescence, and use	Purity	Protein A or G purified
Description CoraFluor(TM) 1 is a high performance terbium-based TR-FRET (Time-Resolved Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1. Does used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and protein quantification assays. Host Mouse Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422) Notes CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 20	Buffer	PBS
Fluorescence Resonance Energy Transfer) or TRF (Time-Resolved Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs 2UV light at approximately 340 nm, and emits at approximately 490 nm 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 to absorb at the emission wavelengths of CoraFluor (TM) 1. CoraFluor(TM) 1. CoraFluor 1. Cor	Product Description	
Gene ID 4072 Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422) Notes CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry/	Description	Fluorescence) donor for high throughput assay development. CoraFluor(IM) 1 absorbs UV light at approximately 340 nm, and emits at approximately 490 nm, 545 nm, 585 nm and 620 nm. It is compatible with common acceptor dyes that absorb at the emission wavelengths of CoraFluor(TM) 1. CoraFluor(TM) 1 can be used for the development of robust and scalable TR-FRET binding assays such as target engagement, ternary complex, protein-protein interaction and
Gene Symbol EPCAM Species Human, Mouse (Negative), Rat (Negative) Reactivity Notes Does not react with Mouse or Rat. Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422) Notes CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/	Host	Mouse
SpeciesHuman, Mouse (Negative), Rat (Negative)Reactivity NotesDoes not react with Mouse or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia ad vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenRecombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422)NotesCoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry. Immunocytochemistry, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry. Immunocytochemistry/	Gene ID	4072
Reactivity NotesDoes not react with Mouse or Rat.MarkerEpithelial MarkerSpecificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenRecombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422)NotesCoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunofluorescence	Gene Symbol	EPCAM
Marker Epithelial Marker Specificity/Sensitivity EGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Immunogen Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422) CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunocytochemistry, Immunofluorescence	Species	Human, Mouse (Negative), Rat (Negative)
Specificity/SensitivityEGP40 is a 40-43kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma.ImmunogenRecombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422)NotesCoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry. Immunohistochemistry. Immunocytochemistry, Immunocytochemistry/	Reactivity Notes	Does not react with Mouse or Rat.
epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep- CAM). It is expressed on baso-lateral cell surface in most simple epithelia and vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. Thi 	Marker	Epithelial Marker
Notes CoraFluor (TM) is a trademark of Bio-Techne Corp. Sold for research purposes only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry, Immunocytochemistry/	Specificity/Sensitivity	CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from
only under agreement from Massachusetts General Hospital. US patent 2022/0025254 Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/	Immunogen	Recombinant full-length human EpCAM/TROP1 protein (Uniprot: P16422)
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/	Notes	
Immunohistochemistry, Immunohistochemistry-Paraffin, ImmunofluorescenceRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/	Product Application Details	
	Applications	
	Recommended Dilutions	



Application Notes	Optimal dilution of this antibody should be experimentally determined.
-------------------	--

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-11478CL1

NBP2-52190-0.05mg	Recombinant Human EpCAM/TROP1 His Protein
202-IL-010	IL-2 [Unconjugated]
960-EP-050	EpCAM/TROP1 [Unconjugated]
AF748	E-Cadherin Antibody [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11478CL1

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

