## **Product Datasheet**

### CD79A Antibody (IGA/764) [DyLight 755] NBP3-11494IR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11494IR

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11494IR



#### NBP3-11494IR

CD79A Antibody (IGA/764) [DyLight 755]

| 0.1 ml   |
|--|
| Please see the vial label for concentration. If unlisted please contact technical services.  |
| Store at 4C in the dark.   |
| Monoclonal   |
| IGA/764  |
| 0.05% Sodium Azide   |
| IgG1 Kappa   |
| DyLight 755  |
| Protein A or G purified  |
| 50mM Sodium Borate   |
|  |
| Mouse  |
| 973  |
| CD79A  |
| Human  |
| B Cell Marker  |
| A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. |
| Recombinant full-length human CD79A protein (Uniprot: P11912)  |
|  |
| DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.  |
|  |
|  |
| Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence,<br>Immunohistochemistry, Immunohistochemistry-Paraffin   |
|  |
|  |





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-11494IR

NBP1-43319IR-0.5mlMouse IgG1 Kappa Isotype Control (P3.6.2.8.1) [DyLight 755]NBP2-60209-50ugRecombinant Human CD79A His Protein7268-CT-100CTLA-4 [Unconjugated]9685-CD-050CD79A [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11494IR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

