Product Datasheet

CD34 Antibody (SPM610) [Alexa Fluor® 488] NBP3-11532AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11532AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11532AF488



NBP3-11532AF488

CD34 Antibody (SPM610) [Alexa Fluor® 488]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM610Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description947Gene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpeciesSpecies a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein on 90-120Ka, which is identified as CD34 (Uirenational workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for stasini gubasis in acute myeloid leukemia. In addition, it is expression is also found in vascular endothelial marker of choice for stasini gubasis in acute myeloid leukemia. In addition, it is expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosaccoma and Kaposi sa scolary.			
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM610Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpeciesThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a halmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelian. Addition, it is expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expression is gradually, poliferating endothelial cells. Avic. CD34 leabels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Product Information		
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneSPM610Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID947Gene SymbolCD34Specificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmebrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells avit-CD34 leabs 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Unit Size	0.1 ml	
ClonalityMonoclonalCloneSPM610Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein or 190-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stronal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpression tis as solitary fibrous tumor and gastrointestinal stronal tumors. Such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibrous tumor and gastrointestinal stronal tumors, such as solitary fibr	Concentration		
CloneSPM610Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as cD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Storage	Store at 4C in the dark.	
Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Clonality	Monoclonal	
IsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endotheliam. Additionally, proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Clone	SPM610	
ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Preservative	0.05% Sodium Azide	
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Isotype	IgG1 Kappa	
Buffer50mM Sodium BorateProduct DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Conjugate	Alexa Fluor 488	
Product DescriptionHostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Purity	Protein A or G purified	
HostMouseGene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Buffer	50mM Sodium Borate	
Gene ID947Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Product Description		
Gene SymbolCD34SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Host	Mouse	
SpeciesHuman, RatMarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Gene ID	947	
MarkerHematopoietic Stem Cell & Endothelial MarkerSpecificity/SensitivityThis antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Gene Symbol	CD34	
Specificity/Sensitivity This antibody recognizes a carbohydrate epitope on a single chain, transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Species	Human, Rat	
transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85% of angiosarcoma and Kaposi s sarcoma, but shows low specificity.	Marker	Hematopoietic Stem Cell & Endothelial Marker	
Immunogen Recombinant full-length human CD34 protein (Uniprot: P28906)	Specificity/Sensitivity	transmembrane, heavily glycosylated protein of 90-120kDa, which is identified as CD34 (VI international workshop on human differentiation antigens). Its expression is a hallmark for identifying pluripotent hematopoietic stem or progenitor cells. Its expression is gradually lost as lineage committed progenitors differentiate. CD34 is a marker of choice for staining blasts in acute myeloid leukemia. In addition, it is expressed by soft tissue tumors, such as solitary fibrous tumor and gastrointestinal stromal tumor. CD34 expression is also found in vascular endothelium. Additionally, proliferating endothelial cells overexpress this molecule than the non-proliferating endothelial cells. Anti-CD34 labels 85%	
	Immunogen	Recombinant full-length human CD34 protein (Uniprot: P28906)	

www.novusbio.com



Notes	

Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.

Product Application Details	
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Images

CD34 Antibody (SPM610) [Alexa Fluor® 488]

	Alexa Fluc	or® 488	
Same and the second	LASER (nm)	FILTER	
Alexa Fluor ^{® 488}	Blue (488)	525/50	
	EXCITATION MAX (nm)	EMISSION MAX (nm)	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-11532AF488

IC002G	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 488]
NBP2-53044-50ug 210-TA-005	Recombinant Human CD34 His Protein TNF-alpha [Unconjugated]
9655-CD-050	CD34 [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11532AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

