Product Datasheet

S100A9 Antibody (S100A9/1075) [Biotin] NBP3-11559B

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11559B

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-11559B



NBP3-11559B

Product Information	S100A9 Antibody (S100A9/1075) [Biotin]		
Concentration Please see the vial tabel for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone \$100A9/1075 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Bymbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupflers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stormach. S100A9 is also expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and enpithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotxok shock in response to bacterial lipopolysacchanide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of ILB and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry-Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry.	Product Information		
Storage Store at 4C in the dark. Clonality Monoclonal Clone S100A9/1075 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specifficity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mante zone B-ymphocytes of the lymn hode and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyce migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence, Immunohistochemistry, Immunophistochemistry, Immunohistochemistry, Immunofluorescence, Immunohistochemistry, Immunohis	Unit Size	0.1 ml	
Clone S100A9/1075 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in a cutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is a cyresed so resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of ILB and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofiuorescence, Immunofiuorescence, Im	Concentration	•	
Clone S100A9/1075 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and fungil. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa B. S100A9 lays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Storage	Store at 4C in the dark.	
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Symbol \$100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiccytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. \$100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. \$100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. \$100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa B. \$100A9 lays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human \$100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Clonality	Monoclonal	
IgG1 Kappa IgG1 Kappa Biotin	Clone	S100A9/1075	
Conjugate Biotin Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marcophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dematoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial plopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of ILS and cell-surface expression of ICAM1. Immunogen Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer PBS Product Description Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-infilammatory mediator in acute and chronic inflammation made up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohi	Isotype	IgG1 Kappa	
Buffer PBS	Conjugate	Biotin	
Product Description Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoletic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence, Immunofluorescence	Purity	Protein A or G purified	
Host Mouse Gene ID 6280 Gene Symbol S100A9 Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermactoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Buffer	PBS	
Gene Symbol Species Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Product Description		
Secies Human, Rat Marker Macrophage Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Host	Mouse	
Species Human, Rat	Gene ID	6280	
Marker Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermaces. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Gene Symbol	S100A9	
Specificity/Sensitivity This monoclonal antibody stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. \$100A9\$ is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. \$100A9\$ is also expressed in epithelial cells constitutively or induced during dermatoses. \$100A9\$ is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. \$100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human \$100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Flow Cytometry, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Species	Human, Rat	
in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1. Immunogen Recombinant human S100A9 protein (Uniprot: P06702) Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry-Paraffin, Immunofluorescence	Marker	Macrophage Marker	
Product Application Details Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence		in hematopoietic organs, Kupffers cells of the liver and Langerhans cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.	
Applications Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence Recommended Dilutions Flow Cytometry, Immunohistochemistry-Paraffin, Immunocytochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	Immunogen	Recombinant human S100A9 protein (Uniprot: P06702)	
Recommended Dilutions Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence			
Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence		Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence	
Application Notes Optimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin,	
	Application Notes	Optimal dilution of this antibody should be experimentally determined.	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-11559B

IC002B Mouse IgG1 Isotype Control (11711) [Biotin]
NBP1-44500 Recombinant Human S100A9 His Protein

210-TA-005 TNF-alpha [Unconjugated]

DY5578 S100A9 [Biotin]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11559B

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

