Product Datasheet

Cytokeratin 8/18 Antibody (C-51) [Alexa Fluor® 488] NBP3-11565AF488

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11565AF488

Updated 10/26/2023 v.20.1

Earn rewards for product reviews and publications.



Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11565AF488

NBP3-11565AF488

Cytokeratin 8/18 Antibody (C-51) [Alexa Fluor® 488]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clonality Monoclonal Clone C-51 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Venepus (Negative), Mouse (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. > Molecular weight cytokeratins and exists in combination with cytokeratin 18 (Crkeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (Crkeratin 18) (Chkeratin 28) carcinomas are usually negative. This antibody is useful in ademocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This sintbody is useful in democar		·	
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneC-51Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Product Information		
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone C-51 Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Unit Size	0.1 ml	
ClonalityMonoclonalCloneC-51Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionMouseHostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Concentration		
CloneC-51Preservative0.05% Sodium AzideIsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. shr/>shr/>rkat reactivity reported in scientific liteRature (PMID: 28975767).Specificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal I tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Storage	Store at 4C in the dark.	
Preservative 0.05% Sodium Azide Isotype IgG1 Kappa Conjugate Alexa Fluor 488 Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >cbr/>cbr/>Spt/>Sptice and antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Clonality	Monoclonal	
IsotypeIgG1 KappaConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >cbr/> >cbr/> >ktr8.Specificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratin and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Clone	C-51	
ConjugateAlexa Fluor 488PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Preservative	0.05% Sodium Azide	
Purity Protein A or G purified Buffer 50mM Sodium Borate Product Description Mouse Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Isotype	IgG1 Kappa	
Buffer 50mM Sodium Borate Product Description Mouse Host Mouse Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >chr/>chr/>Rat reactivity reported in scientific liteRature (PMID: 28975767). Specificity/Sensitivity Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Conjugate	Alexa Fluor 488	
Product DescriptionHostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >kpr/> Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Purity	Protein A or G purified	
HostMouseGene ID3856Gene SymbolKRT8SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >kpr/> Specificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Buffer	50mM Sodium Borate	
Gene ID 3856 Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >br/> >kpr/>Rat reactivity reported in scientific liteRature (PMID: 28975767). Specificity/Sensitivity Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Product Description		
Gene Symbol KRT8 Species Human, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative) Reactivity Notes Does not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. 	Host	Mouse	
SpeciesHuman, Rat, Porcine, Bovine, Ferret (Negative), Monkey, Sheep, Chicken (Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. >kpr/> Specificity/SensitivityDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. (Vegative)Specificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Gene ID	3856	
(Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus (Negative)Reactivity NotesDoes not react with Mouse, Ferret, Rabbit, donkey, Chicken or Xenopus laevis. Specificity/SensitivitySpecificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Gene Symbol	KRT8	
 Specificity/SensitivityCytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Species	(Negative), Donkey (Negative), Mouse (Negative), Rabbit (Negative), Xenopus	
molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the	Reactivity Notes		
normal epidermis so only Paget cells are stained.	Specificity/Sensitivity	molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This monoclonal antibody recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in	
Immunogen Cytoskeleton preparation from HeLa cells	Immunogen	Cytoskeleton preparation from HeLa cells	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.	
Product Application Details		
Applications	Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Immunofluorescence	
Recommended Dilutions	Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Immunofluorescence	
Application Notes	Optimal dilution of this antibody should be experimentally determined.	

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-11565AF488

MAB1368	alpha-Fetoprotein/AFP Antibody (189502) [Unconjugated]
NB300-223	Vimentin Antibody
8184-CK-050	Choline Kinase beta [Unconjugated]
IC002G	Mouse IgG1 Isotype Control (11711) [Alexa Fluor® 488]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11565AF488

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

