Product Datasheet

CD79A Antibody (IGA/1790R) [Alexa Fluor® 405] NBP3-11617AF405

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11617AF405

Updated 11/12/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11617AF405



NBP3-11617AF405

CD79A Antibody (IGA/1790R) [Alexa Fluor® 405]

| Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIGA/1790RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionFluor 405HostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-inked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on a tracellular component. CD79a is tage pars at pre B cell istage, early in maturation, and persists until the plasma cell istage where it is found as an intracellular component. CD79a is touch in the majority of acute level as an intracellular component. CD79a is sound in the majority of acute level as anti-CD20. Nut also is more likely to stain B-1/mphomasa, and is one myelomas. It is not present in myeloid or T cell lines, B cell stage were it is found as an intracellular component. CD79a is sound in the majority of acute levelmans de norcellorent minutor200, This antibody will stain many of the | | |
|---|-------------------------|--|
| ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIGA/1790RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct Description973HostRabbitGene BD973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells | Product Information | |
| services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneIGA/1790RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Ag receptor. CD79a inst appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precurs or Beerli Type, in G cell Ising, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precurs or Beerli Type, in G cell Ising, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precurs or Beerli Type, in G cell Ising, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precurs or Beerli Type, in G cell Ising, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the same lymphomas aterImmunogenA synthetic pe | Unit Size | 0.1 ml |
| ClonalityMonoclonalCloneIGA/1790RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cell Ag receptor. CD79a first appears at pre B cell stage early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, Anti-CD79a is generally used to complement anti-CD20. This antiboddy will stain many of the same lymphomas. It is not present in myeloid or T cell lines, and cocasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Concentration | • |
| CloneIGA/1790RPreservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mot-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin compre | Storage | Store at 4C in the dark. |
| Preservative0.05% Sodium AzideIsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polype | Clonality | Monoclonal |
| IsotypeIgGConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Ntils or mature B-cell stage merel bymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Clone | IGA/1790R |
| ConjugateAlexa Fluor 405PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypeptides and immunoglobulins on B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Preservative | 0.05% Sodium Azide |
| PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cell Stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines, Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Isotype | IgG |
| Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins constitute the B cell Agr ecceptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomaleukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Conjugate | Alexa Fluor 405 |
| Product DescriptionHostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is gound in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/eukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Purity | Protein A purified |
| HostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/eukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Buffer | 50mM Sodium Borate |
| Gene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Product Description | |
| Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Host | Rabbit |
| SpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Gene ID | 973 |
| Reactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Gene Symbol | CD79A |
| MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Species | Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey |
| Specificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of | Reactivity Notes | Others not known. |
| CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen | Marker | B-Cell Marker |
| | Specificity/Sensitivity | CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. |
| | Immunogen | |

www.novusbio.com



| | Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet. |
|-----------------------------|---|
| Product Application Details | |
| •• | |
| Applications | Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array |
| Recommended Dilutions | Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Protein Array |
| Application Notes | Optimal dilution of this antibody should be experimentally determined. |
| 1 | |

Notes





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-11617AF405

| NBP2-24891AF405 | Rabbit IgG Isotype Control [Alexa Fluor® 405] |
|-----------------|---|
| NBP2-60209-50ug | Recombinant Human CD79A His Protein |
| 7268-CT-100 | CTLA-4 [Unconjugated] |
| 9685-CD-050 | CD79A [Unconjugated] |
| | |

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11617AF405

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

