Product Datasheet

CD79A Antibody (IGA/1790R) [DyLight 755] NBP3-11617IR

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-11617IR

Updated 11/12/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-11617IR



NBP3-11617IR

CD79A Antibody (IGA/1790R) [DyLight 755]

Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone IGA/1790R Preservative 0.05% Sodium Azide Isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity Gell Marker Specificity/Sensitivity Gore Sourge of meature of presense of the presense of the specificity of a date is found as an intracellular component. CD79a is found in the majority of acute is found as an intracellular component. CD79a is found in the majority of acute is found as an intracellular component. CD79a is found in the majority of acute is found as an intracellular coresponding via as sotaris action of the specificity of a also staim smary. Specificity/Sensitivity Mothoccanalition and accasionaliy some types of endothelial	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone IGA/1790R Preservative 0.05% Sodium Azide Isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity CD79A Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulin constitute the B cell Agreecptor. CD79a inst appears at pre B cell stage, early in maturation, and persists until the plasma cell asge where it is found as an intracellular component. CD79a is squerally used to complement anti-CD20. but also is more likely to stains more cases of plasma cell myrelom as a coll myrelof or T cell lines, Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stains more cases of plasma cell myrelom as a coll myrelof or T cell lines, So tains more cases of plasma cell myrelom as a coll set set ocomplement anti-CD20. Dut also is more likely to stains more cases of plasma cell myrelomas	Unit Size	0.1 ml
Clonality Monoclonal Clone IGA/1790R Preservative 0.05% Sodium Azide Isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Host Rabbit Gene D 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and every corcers of cell type, in B cell intes, B cell stage where it is found as an intracellular component. CD79a is generally used to complement anti-CD20. Jut is o is more likely to statis by mphomas atter treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomaleukemias of precursor B cell type, in B cell imphomas atter Immunogen A synthetic peptide corresponding to as 20:216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark	Concentration	•
Clone IGA/1790R Preservative 0.05% Sodium Azide Isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer Som Sodium Borate Product Description Host Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD78b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 oplypenpides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypenpides and immunoglobulins on B cells. This complex of mb-1 and B29 oplypenpides and immunoglobulins on B cells. This complex sturil the plasma cell stage were it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell Inset. B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Ant-CD78a is generally used to complement anti-CD20. Nut -LCD79a atos stains more cases of plasma cell myeloma and anti-CD20. Jut also is more likely to stain. B-lymphoblastic tymphoma% as anti-CD20. Nut -LCD79a atos stains more cases of plasma cell myeloma and anti-CD20. Jut also is more likely to stain. B-lymphoblastic Notes <t< th=""><th>Storage</th><th>Store at 4C in the dark.</th></t<>	Storage	Store at 4C in the dark.
Preservative 0.05% Sodium Azide Isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides. Sound as an intracellular component. CD79a is found in the mapiority of acutte leukemias ant-CD20. Usepecially for ma	Clonality	Monoclonal
isotype IgG Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Image: Conjugate Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukernias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell Times. Anti-CD79a is generally unsert to complement anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark	Clone	IGA/1790R
Conjugate DyLight 755 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypecifies and immunoglobulins on stitute the B cell Ag receptor. CD79a (ist appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. Anti-CD79a is generally used to complement anti-CD20 but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/eukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and coccasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to as 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product	Preservative	0.05% Sodium Azide
Purity Protein A purified Buffer 50mM Sodium Borate Product Description Rabbit Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cell Stage. Early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD29 is generally used to complement anti-CD20. but also is more likely to stain B-lympholastic lymphomas as anti-CD20, but also is more likely to stain B-lympholastic lymphoma/eukemia than is anti-CD20. Anti-CD79 al aso stains more cases of plasma cell myeloma ant occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to a 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.	Isotype	IgG
Buffer 50mM Sodium Borate Product Description Rabbit Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. Stain B-lymphoblastic lymphomas atter treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoman/cukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to a 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Prod	Conjugate	DyLight 755
Product Description Host Rabbit Gene ID 973 Gene Symbol CD79A Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on a scale. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. A nti-CD79a is generally used to complement anti-CD20 operais found in the majority of acute leukemias of precursor B cell type, in B cell lines. A nti-CD79a is generally used to complement anti-CD20. Ut also is more likely to stain B-tymphomas after treatment with Rituximab (anti-CD20). Ut also is more likely to stain B-tympholastic lymphomas as anti-CD20. Ut also is more likely to stain B-tympholastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) </th <th>Purity</th> <th>Protein A purified</th>	Purity	Protein A purified
HostRabbitGene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in some myelomas. It is not present in myeloid or T cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to a 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application Details ApplicationsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Protein Array	Buffer	50mM Sodium Borate
Gene ID973Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell stage energily used to complement anti-CD20. Nati-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Paraffin, Protein Array	Product Description	
Gene SymbolCD79ASpeciesHuman, Mouse, Rat, Porcine, Bovine, Canine, MonkeyReactivity NotesOthers not known.MarkerB-Cell MarkerSpecificity/SensitivityA disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mc1 and B29 polypeptides and immunoglobulin son B cells. This complex of mc1 and B29 users at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphomas and corcasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein ArrayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Paraffin, Protein Array	Host	Rabbit
Species Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same hymphomas as anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Immunochistochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Paraffin, Protein Array	Gene ID	973
Reactivity Notes Others not known. Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry-Immunofluorescence, Immunohistochemistry, Paraffin, Protein Array	Gene Symbol	CD79A
Marker B-Cell Marker Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry-Paraffin, Protein Array	Species	Human, Mouse, Rat, Porcine, Bovine, Canine, Monkey
Specificity/Sensitivity A disulphide-linked heterodimer, consisting of mb-1 (or CD79a) and B29 (or CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines. A cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20. but also is more likely to stain B-lymphoblastic lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well. Immunogen A synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein Array	Reactivity Notes	Others not known.
CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 bepecially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of plasma cell myeloma and occasionally some types of endothelial cells as well.ImmunogenA synthetic peptide corresponding to aa 202-216 (GTYQDVGSLNIADVQ) of human CD79A protein. (Uniprot: P11912)NotesDyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.Product Application DetailsWestern Blot, Flow Cytometry, Immunohistochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein Array	Marker	B-Cell Marker
human CD79A protein. (Uniprot: P11912) Notes DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Immunohistochemistry, Paraffin, Protein Array	Specificity/Sensitivity	CD79b) polypeptides, is non-covalently associated with membrane-bound immunoglobulins on B cells. This complex of mb-1 and B29 polypeptides and immunoglobulin constitute the B cell Ag receptor. CD79a first appears at pre B cell stage, early in maturation, and persists until the plasma cell stage where it is found as an intracellular component. CD79a is found in the majority of acute leukemias of precursor B cell type, in B cell lines, B cell lymphomas, and in some myelomas. It is not present in myeloid or T cell lines. Anti-CD79a is generally used to complement anti-CD20 especially for mature B-cell lymphomas after treatment with Rituximab (anti-CD20). This antibody will stain many of the same lymphomas as anti-CD20, but also is more likely to stain B-lymphoblastic lymphoma/leukemia than is anti-CD20. Anti-CD79a also stains more cases of
DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. Product Application Details Applications Western Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunohis	Immunogen	
ApplicationsWestern Blot, Flow Cytometry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry, Immunohistochemistry-Paraffin, Protein ArrayRecommended DilutionsWestern Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Protein Array	Notes	DyLight (R) is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.
Immunohistochemistry, Immunohistochemistry-Paraffin, Protein Array Recommended Dilutions Western Blot, Flow Cytometry, Immunohistochemistry, Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Protein Array	Product Application Details	
Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin, Protein Array	Applications	
Application NotesOptimal dilution of this antibody should be experimentally determined.	Recommended Dilutions	Immunocytochemistry/Immunofluorescence, Immunohistochemistry-Paraffin,
	Application Notes	Optimal dilution of this antibody should be experimentally determined.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-11617IR

NBP2-24891IRRabbit IgG Isotype Control [DyLight 755]NBP2-60209-50ugRecombinant Human CD79A His Protein7268-CT-100CTLA-4 [Unconjugated]9685-CD-050CD79A [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-11617IR

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

