Product Datasheet

Fc epsilon RI Antibody (IB10) - Chimeric - Azide and BSA Free NBP3-12050-0.2mg

Unit Size: 0.2 mg

Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-12050

Updated 4/19/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications
Submit a review at www.novusbio.com/reviews/destination/NBP3-12050



NBP3-12050-0.2mg

Fc epsilon RI Antibody (IB10) - Chimeric - Azide and BSA Free


, ,		
Product Information		
Unit Size	0.2 mg	
Concentration	1 mg/ml	
Storage	Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.	
Clonality	Monoclonal	
Clone	IB10	
Preservative	0.02% Proclin 300	
Isotype	IgG Kappa	
Purity	Protein A purified	
Buffer	PBS	

Product Description	
Host	Rabbit
Gene ID	2205
Gene Symbol	FCER1A
Species	Human
Specificity/Sensitivity	This antibody binds to human Fc epsilon receptor and competes with IgE binding.
Immunogen	Recombinant human protein, N-terminus.

Product Application Details	
Applications	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Block/Neutralize
Recommended Dilutions	Flow Cytometry, Immunocytochemistry/Immunofluorescence, Block/Neutralize
Application Notes	IB10 competes with IgE for binding to human FCER1A; consequently, its binding to mast cells and basophils is decreased when IgE concentration is high, but it does block IgE binding to these cells. IB10 and IE7 do not crossblock each other's binding. This chimeric rabbit antibody was made using the variable domain sequences of the original Mouse IgG1 format, for improved compatibility with existing reagents, assays and techniques.

Images

Immunocytochemistry/Immunofluorescence: Fc epsilon RI Antibody (IB10) - Chimeric [NBP3-12050] - Immunofluorescence analysis of paraformaldehyde fixed U937 cells immobilized on Shi-fix(TM) coverslips and stained with the chimeric rabbit IgG version of IB10 (NBP3-12050) at 10 ug/ml followed by Alexa Fluor(R) 488 secondary antibody (2 ug/ml), showing diffuse membrane staining. The nuclear stain is DAPI (blue). Panels show from left-right, top-bottom NBP3-12050, DAPI, merged channels and an isotype control. The isotype control was stained with anti-Fluorescein antibody (NBP2-52638) followed by Alexa Fluor(R) 488 secondary antibody.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966 novus@novusbio.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom

Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: technical@novusbio.com

Orders: orders@novusbio.com General: novus@novusbio.com

Products Related to NBP3-12050-0.2mg

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7156 Goat anti-Rabbit IgG (H+L) Secondary Antibody
NBP2-23043 Recombinant Human Fc epsilon RI His Protein

210-TA-005 TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-12050

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

