## **Product Datasheet**

### Decorin Antibody (DCN/7031R) [Alexa Fluor® 532] NBP3-14026AF532

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-14026AF532

Updated 10/26/2023 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-14026AF532



#### NBP3-14026AF532

Decorin Antibody (DCN/7031R) [Alexa Fluor® 532]

Product InformationUnit Size0.1 mlConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneDCN/7031RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1634Gene ID1634Gene SymbolDCNSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains the formation of fibers with variations in stability and solubility. The Decorin binds to Collagen Type I, Il and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin binds to growt factors, intercellular matrix molecules, such as Fibornection or fibers with variations in stability and solubility. The Decorin binds to and inhibits TGF may contribute to cartilage destruction that is Fibornection or the optical startic of inflammatory joint diseases. For immunostaining pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact sequence is proprietary) (Uniprot: PO7S85)		-
ConcentrationPlease see the vial label for concentration. If unlisted please contact technical services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneDCN/7031RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1634Gene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains is the lug and soliding bord stabilizing loops. Decorin binds to Gollagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to Grow factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chordroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Product Information	
services.StorageStore at 4C in the dark.ClonalityMonoclonalCloneDCN/7031RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct Description1634Gene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminog flycan chain-containing core protein. The core protein contains the neucine rich repeats that contain sites for glycosylation, flanked by disulfide bord stabilizing loops. Decorin binds to Gollagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to Growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to Growth factors, intercellular matrix molecules, such as Fibronectin of fibers may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chordroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Unit Size	0.1 ml
ClonalityMonoclonalCloneDCN/7031RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionKabbitHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpeciesHumanSpeciesburden containing core protein. The core protein contains ten leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability. The Decorin core protein and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin and to the Decorin endocytosis receptor. Decorin binds to G or inflammatory joint diseases. For immunostaining, pre-incubation with chondrotitnase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Concentration	•
CloneDCN/7031RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpeciesHumanSpeciesSocial containing core protein. The core protein contains test for a dycosplan, and the formation of fibers with variations in stability and solubility. The Decorin binds to got matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bord stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Clonality	Monoclonal
IsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Clone	DCN/7031R
ConjugateAlexa Fluor 532PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Preservative	0.05% Sodium Azide
PurityProtein A or G purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Isotype	IgG Kappa
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Conjugate	Alexa Fluor 532
Product DescriptionHostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Purity	Protein A or G purified
HostRabbitGene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Buffer	50mM Sodium Borate
Gene ID1634Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Product Description	
Gene SymbolDCNSpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Host	Rabbit
SpeciesHumanSpecificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Gene ID	1634
Specificity/SensitivityDecorin is a small leucine-rich proteoglycan (SLRP) family member that consists of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Gene Symbol	DCN
of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.ImmunogenRecombinant human Decorin protein fragment (around aa212-336) (exact	Species	Human
	Specificity/Sensitivity	of a glycosaminoglycan chain-containing core protein. The core protein contains ten leucine rich repeats that contain sites for glycosylation, flanked by disulfide bond stabilizing loops. Decorin binds to Collagen Type I, II and IV in vivo and promotes the formation of fibers with variations in stability and solubility. The Decorin core protein binds to growth factors, intercellular matrix molecules, such as Fibronectin and Thrombospondin, and to the Decorin endocytosis receptor. Decorin binds to and inhibits TGF may contribute to cartilage destruction that is characteristic of inflammatory joint diseases. For immunostaining, pre-incubation with chondroitinase-SBC or testicular hyaluronidase may be required to expose the epitope.
	Immunogen	

www.novusbio.com



	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Immunohistochemistry-Paraffin
<b>Recommended Dilutions</b>	Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.

Notes





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 novus@novusbio.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: technical@novusbio.com Orders: orders@novusbio.com General: novus@novusbio.com

#### Products Related to NBP3-14026AF532

NBP1-91258	Fibronectin Antibody - BSA Free
AF808	Osteopontin/OPN Antibody [Unconjugated]
DY143	Decorin [Biotin]
NBP2-56346PEP	Decorin Recombinant Protein Antigen

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-14026AF532

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

