# **Product Datasheet**

# Biotin (Vitamin B7 or Vitamin H) Antibody (BTN/403) -Azide and BSA Free NBP3-14087

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-14087

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-14087

#### NBP3-14087

Biotin (Vitamin B7 or Vitamin H) Antibody (BTN/403) - Azide and BSA Free

100 ug	
1 mg/ml	
Store at -20 to -80C. Avoid freeze-thaw cycles.	
Monoclonal	
BTN/403	
No Preservative	
IgG1 Kappa	
Protein A or G purified	
10mM PBS	
Product Description	
<ul> <li>1.0 mg/ml of antibody purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS WITHOUT BSA &amp; azide. Also available at 200 ug/ml WITH BSA &amp; azide (NBP3-13904).</li> <li>Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80 C.</li> </ul>	
Mouse	
Non-species specific	
It recognizes both the free and protein-conjugated (either soluble or cell bound) form of biotin. This monoclonal antibody is highly specific to biotin and shows no cross-reaction with other structurally related compounds. It has a very high affinity for biotin and is excellent for use in various amplification techniques. In some applications, localization of biotinylated probes with avidin produces unacceptably high background staining. Anti-biotin antibody may be substituted to decrease this noise.	
Biotinylated sheep immunoglobulin	
Product Application Details	
Western Blot, Flow Cytometry, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry-Paraffin, In-situ Hybridization	
Western Blot 0.5-2 ug/ml, Flow Cytometry 0.5-1 ug/million cells, Immunocytochemistry/ Immunofluorescence 0.5-1 ug/ml, Immunohistochemistry- Paraffin 0.5-1 ug/ml, In-situ Hybridization	
Confocal Microscopy Immunohistochemistry (Formalin-fixed): 0.5-1ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes In situ Nucleic Acid Hybridization	





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

## **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

## **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

# Products Related to NBP3-14087

HAF007	Goat anti-Mouse IgG Secondary Antibody [HRP]
NB720-B	Rabbit anti-Mouse IgG (H+L) Secondary Antibody [Biotin]
NBP1-43319-0.5mg	Mouse IgG1 Kappa Isotype Control (P3.6.2.8.1)

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-14087

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

