# **Product Datasheet**

# alpha-Sarcoglycan Antibody (3W3Q10) NBP3-15896-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-15896

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-15896



#### NBP3-15896-100ul

alpha-Sarcoglycan Antibody (3W3Q10)


| Product Information |   |
|---------------------|---|
| Unit Size           | 100 ul  |
| Concentration       | Please see the vial label for concentration. If unlisted please contact technical services. |
| Storage             | Store at -20C. Avoid freeze-thaw cycles.  |
| Clonality           | Monoclonal  |
| Clone               | 3W3Q10  |
| Preservative        | 0.02% Sodium Azide  |
| Isotype             | IgG   |
| Purity              | Affinity purified   |
| Buffer              | PBS (pH 7.3), 50% glycerol, 0.05% BSA   |
| Product Description |   |

| Product Description |  |
|---------------------|--|
| Description         | Novus Biologicals Rabbit alpha-Sarcoglycan Antibody (3W3Q10) (NBP3-15896) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.  |
| Host                | Rabbit   |
| Gene ID             | 6442   |
| Gene Symbol         | SGCA   |
| Species             | Mouse, Rat   |
| Immunogen           | A synthetic peptide corresponding to a sequence within amino acids 288-387 of human alpha-Sarcoglycan (SGCA) (Q16586). VDALVTLLVPLLVALLLTLLLAYVMCCRREGRLKRDLATSDIQMVHHCTIHGNTE ELRQMAASREVPRPLSTLPMFNVHTGERLPPRVDSAQVPLILDQH |

| <b>Product Application Details</b> |  |
|------------------------------------|--|
| Applications                       | Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/<br>Immunofluorescence, Immunohistochemistry                                      |
|                                    | Western Blot 1:100 - 1:500, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 1:50 - 1:200, Immunohistochemistry-Paraffin |

# **Images**


Western Blot: alpha-Sarcoglycan Antibody (3W3Q10) [NBP3-15896] - Western blot analysis of extracts of various cell lines, using alpha-Sarcoglycan (SGCA) Rabbit mAb (NBP3-15896) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.





Page 2 of 3 v.20.1 Updated 9/9/2025 Immunohistochemistry-Paraffin: alpha-Sarcoglycan Antibody (3W3Q10) [NBP3-15896] - Immunohistochemistry of paraffin-embedded mouse heart using alpha-Sarcoglycan (SGCA) Rabbit mAb (NBP3-15896) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. Immunohistochemistry-Paraffin: alpha-Sarcoglycan Antibody (3W3Q10) [NBP3-15896] - Immunohistochemistry of paraffin-embedded rat heart using alpha-Sarcoglycan (SGCA) Rabbit mAb (NBP3-15896) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol. alpha Sarcoglycan(SGCA) Immunocytochemistry/ Immunofluorescence: alpha-Sarcoglycan Antibody (3W3Q10) [alpha-Sarcoglycan] - Confocal imaging of paraffinembedded Rat heart using alpha-Sarcoglycan Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining. Immunocytochemistry/ Immunofluorescence: alpha-Sarcoglycan Antibody (3W3Q10) [alpha-Sarcoglycan] - Confocal imaging of paraffinembedded Mouse heart using alpha-Sarcoglycan Rabbit mAb followed

by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L). DAPI was used for nuclear staining (Blue). Objective: 40x. Perform high pressure antigen retrieval with 0.01 M citrate buffer (pH 6.0) prior to IF staining.





# Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

### Products Related to NBP3-15896-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

H00006442-P01-10ug Recombinant Human alpha-Sarcoglycan GST (N-Term) Protein

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-15896

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

