Product Datasheet

Prohibitin 2 Antibody (8I10B1) NBP3-16766-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-16766

Updated 9/9/2025 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-16766



NBP3-16766-100ul

Prohibitin 2 Antibody (8I10B1)


Product Information	
Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	8I10B1
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol, 0.05% BSA
Product Description	

Product Description	
Description	Novus Biologicals Rabbit Prohibitin 2 Antibody (8I10B1) (NBP3-16766) is a recombinant monoclonal antibody validated for use in IHC, WB, ICC/IF and IP. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	11331
Gene Symbol	PHB2
Species	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human Prohibitin 2 (Prohibitin 2) (Q99623). QMVNISLRVLSRPNAQELPSMYQRLGLDYEERVLPSIVNEVLKSVVAKFNASQL ITQRAQVSLIRRELTERAKDFSLILDDVAITELSFSREYTAAVEAK


Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation
Recommended Dilutions	Western Blot 1:3000 - 1:6000, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 1:50 - 1:200, Immunoprecipitation 1:500 - 1:1000, Immunohistochemistry-Paraffin

Images


Western Blot: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Western blot analysis of extracts of various cell lines, using Prohibitin 2 (Prohibitin 2) Rabbit mAb (NBP3-16766) at 1:5000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.




Immunocytochemistry/Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Immunofluorescence analysis of U-2 OS cells using Prohibitin 2 (Prohibitin 2) Rabbit mAb (NBP3-16766) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry-Paraffin: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Immunohistochemistry of paraffin-embedded human colon carcinoma using Prohibitin 2 (Prohibitin 2) Rabbit mAb (NBP3-16766) at dilution of 1:100 (40x lens).Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with IHC staining protocol.




Immunocytochemistry/Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Immunofluorescence analysis of NIH-3T3 cells using Prohibitin 2 (Prohibitin 2) Rabbit mAb (NBP3-16766) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunoprecipitation: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Immunoprecipitation analysis of 300ug extracts of SKOV3 cells using 3ug Prohibitin 2 antibody (NBP3-16766). Western blot was performed from the immunoprecipitate using Prohibitin 2 antibody (NBP3-16766) at a dilition of 1:1000.




Immunocytochemistry/ Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Confocal imaging of NIH/3T3 cells using Prohibitin 2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) . The cells were counterstained with alpha-Tubulin Mouse mAb followed by incubation with ABflo(R) 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.




Immunocytochemistry/ Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [NBP3-16766] - Confocal imaging of U-2 OS cells using Prohibitin 2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L) . The cells were counterstained with alpha-Tubulin Mouse mAb followed by incubation with ABflo(R) 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunocytochemistry/ Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [Prohibitin 2] - Confocal imaging of NIH/3T3 cells using Prohibitin 2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L). The cells were counterstained with alpha-Tubulin Mouse mAb followed by incubation with ABflo 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.



Immunocytochemistry/ Immunofluorescence: Prohibitin 2 Antibody (8I10B1) [Prohibitin 2] - Confocal imaging of U-2 OS cells using Prohibitin 2 Rabbit mAb followed by a further incubation with Cy3 Goat Anti-Rabbit IgG (H+L). The cells were counterstained with alpha-Tubulin Mouse mAb followed by incubation with ABflo 488-conjugated Goat Anti-Mouse IgG (H+L) Ab (Green). DAPI was used for nuclear staining (Blue). Objective: 100x.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-16766-100ul

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP2-24891 Rabbit IgG Isotype Control

NBP2-23365 Recombinant Human Prohibitin 2 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-16766

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

