

Product Datasheet

UBE2N/Ubc13 Antibody (1009Z5) NBP3-16793-100ul

Unit Size: 100 ul

Store at -20C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:
www.novusbio.com/NBP3-16793

Updated 9/9/2025 v.20.1

Earn rewards for product
reviews and publications.

Submit a publication at www.novusbio.com/publications

Submit a review at www.novusbio.com/reviews/destination/NBP3-16793



NBP3-16793-100ul

UBE2N/Ubc13 Antibody (10O9Z5)

Product Information

Unit Size	100 ul
Concentration	Please see the vial label for concentration. If unlisted please contact technical services.
Storage	Store at -20C. Avoid freeze-thaw cycles.
Clonality	Monoclonal
Clone	10O9Z5
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Affinity purified
Buffer	PBS (pH 7.3), 50% glycerol, 0.05% BSA

Product Description


Description	Novus Biologicals Rabbit UBE2N/Ubc13 Antibody (10O9Z5) (NBP3-16793) is a recombinant monoclonal antibody validated for use in IHC, WB and ICC/IF. All Novus Biologicals antibodies are covered by our 100% guarantee.
Host	Rabbit
Gene ID	7334
Gene Symbol	UBE2N
Species	Human, Mouse, Rat
Immunogen	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human UBE2N/Ubc13 (P61088). MAGLPRRIKETQRLLAEPVPGIKAEPDESNARYFHVVIAGPQDSPFEGGTFKLE LFLPEEYPMAAPKVRFM TKIYHPNVDKLGRICLDILKDKWSPALQ

Product Application Details

Applications	Western Blot, Immunohistochemistry-Paraffin, Immunocytochemistry/Immunofluorescence, Immunohistochemistry
Recommended Dilutions	Western Blot 1:500 - 1:1000, Immunohistochemistry 1:50 - 1:200, Immunocytochemistry/ Immunofluorescence 1:50 - 1:200, Immunohistochemistry-Paraffin

Images


Western Blot: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Western blot analysis of extracts of various cell lines, using UBE2N/Ubc13 Rabbit mAb (NBP3-16793) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.




Immunocytochemistry/Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Immunofluorescence analysis of U-2 OS cells using UBE2N/Ubc13 Rabbit mAb (NBP3-16793) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry-Paraffin: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Immunohistochemistry of paraffin-embedded human lung cancer using UBE2N/Ubc13 Rabbit mAb (NBP3-16793) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.




Immunocytochemistry/Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Immunofluorescence analysis of NIH-3T3 cells using UBE2N/Ubc13 Rabbit mAb (NBP3-16793) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunocytochemistry/ Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Immunofluorescence analysis of U-2 OS cells using UBE2N/Ubc13 Rabbit mAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunocytochemistry/ Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [NBP3-16793] - Immunofluorescence analysis of NIH/3T3 cells using UBE2N/Ubc13 Rabbit mAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunocytochemistry/ Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [UBE2N/Ubc13] - Immunofluorescence analysis of NIH/3T3 cells using UBE2N/Ubc13 Rabbit mAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunocytochemistry/ Immunofluorescence: UBE2N/Ubc13 Antibody (10O9Z5) [UBE2N/Ubc13] - Immunofluorescence analysis of U-2 OS cells using UBE2N/Ubc13 Rabbit mAb at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.





Novus Biologicals USA

10730 E. Briarwood Avenue
Centennial, CO 80112
USA
Phone: 303.730.1950
Toll Free: 1.888.506.6887
Fax: 303.730.1966
nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave
Toronto, ON M8Z 4E6
Canada
Phone: 905.827.6400
Toll Free: 855.668.8722
Fax: 905.827.6402
canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane
Abingdon Science Park
Abingdon, OX14 3NB, United Kingdom
Phone: (44) (0) 1235 529449
Free Phone: 0800 37 34 15
Fax: (44) (0) 1235 533420
info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com
Technical Support: nb-technical@bio-techne.com
Orders: nb-customerservice@bio-techne.com
General: novus@novusbio.com

Products Related to NBP3-16793-100ul

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP1-51063-0.1mg	Recombinant Human UBE2N/Ubc13 His Protein

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-16793

Earn gift cards/discounts by submitting a publication using this product:
www.novusbio.com/publications

