

# Product Datasheet

## STING/TMEM173 Antibody (2922D) NBP3-18816

Unit Size: 0.1 mg

Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.

[www.novusbio.com](http://www.novusbio.com)



[technical@novusbio.com](mailto:technical@novusbio.com)

Protocols, Publications, Related Products, Reviews, Research Tools and Images at:  
[www.novusbio.com/NBP3-18816](http://www.novusbio.com/NBP3-18816)

Updated 2/3/2025 v.20.1

Earn rewards for product  
reviews and publications.

Submit a publication at [www.novusbio.com/publications](http://www.novusbio.com/publications)

Submit a review at [www.novusbio.com/reviews/destination/NBP3-18816](http://www.novusbio.com/reviews/destination/NBP3-18816)



**NBP3-18816**

STING/TMEM173 Antibody (2922D)


Product Information	
Unit Size	0.1 mg
Concentration	1.0 mg/ml
Storage	Store at 4C for up to 3 months. For longer storage, aliquot and store at -20C.
Clonality	Monoclonal
Clone	2922D
Preservative	0.02% Sodium Azide
Isotype	IgG
Purity	Protein A or G purified
Buffer	PBS

Product Description	
Host	Rabbit
Gene ID	340061
Gene Symbol	STING1
Species	Human, Mouse, Rat
Immunogen	Partial recombinant protein made to amino acids 215-379 of human TMEM173/STING (UniProt Q86WV6).


Product Application Details	
Applications	Western Blot, Simple Western, Immunocytochemistry/ Immunofluorescence, Immunohistochemistry, Immunoprecipitation, Knockout Validated, Multiplex Immunofluorescence
Recommended Dilutions	Western Blot 1 - 2 ug/ml, Simple Western, Immunohistochemistry 0.5 ug/ml, Immunocytochemistry/ Immunofluorescence 3 ug/ml, Immunoprecipitation Validated for Immunoprecipitation from YCharOS Inc. (ycharos.com), Knockout Validated Validated for Knockout from YCharOS Inc. (ycharos.com), Multiplex Immunofluorescence 10 ug/mL

**Images**


STING/TMEM173 was detected in immersion fixed paraffin-embedded sections of human brain Cortex (Cerebrum) using Rabbit Anti-Human STING/TMEM173, Monoclonal Antibody (Catalog #NBP3-18816) at 10ug/mL at 37 ° Celsius for 4 minutes. Before incubation with the primary antibody, tissue underwent an all-in-one dewaxing and antigen retrieval preprocessing using PreTreatment Module (PT Module) and Dewax and HIER Buffer H (pH 9; EpreDia Catalog # TA-999-DHBH). Tissue was stained using the Alexa Fluor™ Plus 647 Goat anti-Rabbit IgG Secondary Antibody at 1:200 at 37 ° Celsius for 2 minutes. (Yellow; Lunaphore Catalog # [DR647RB](#)) and counterstained with DAPI (blue; Lunaphore Catalog # [DR100](#)). Specific staining was localized to the membrane. Protocol available in [COMET™ Panel Builder](#). The image is attached.



**Western Blot: STING/TMEM173 Antibody (2922D) [NBP3-18816]** - Western blot shows cell lysates K562, THP-1, C2C12, and C6. Membrane was probed with 1 ug/mL of (Catalog # NBP3-18816) followed by HRP-conjugated Anti-Rabbit IgG Secondary Antibody (Catalog # HAF008). A specific band was detected for STING/TMEM173 at approximately 42 kDa (as indicated). This experiment was conducted under reducing conditions and using Immunoblot Buffer Group 1. Internally validated Western blot.




**Immunocytochemistry/Immunofluorescence: STING/TMEM173 Antibody (2292D) [NBP3-18816]** - STING/TMEM173 was detected in immersion fixed U937 human myeloid leukaemia cell line but is not detected in Daudi cell line using Rabbit Anti-Human STING/TMEM173 Monoclonal Antibody (Catalog # NBP3-18816) at 1 ug/mL for 3 hours at room temperature. Cells were stained using the NorthernLights(TM) 557-conjugated Anti-Rabbit IgG Secondary Antibody (red; Catalog # NL004) and counterstained with DAPI (blue).




U937 (Positive) cells

Daudi (Negative) cells


**Immunohistochemistry: STING/TMEM173 Antibody (2292D) [NBP3-18816]** - STING/TMEM173 was detected in immersion fixed paraffin-embedded sections of mouse lung tissue using Rabbit Anti-Human STING/TMEM173 Monoclonal Antibody at 0.5 ug/mL for 1 hour at room temperature followed by incubation with the Anti-Rabbit IgG VisUCyte™ HRP Polymer Antibody. Tissue was stained using DAB (brown) and counterstained with hematoxylin (blue). Specific staining was localized to pneumocytes and alveolar cells. Staining with VisUCyte HRP Polymer Detection Reagents.




**Simple Western: STING/TMEM173 Antibody (2922D) [NBP3-18816]** - Simple Western Lane view shows K562 and THP-1 whole cell lysate (WCL). A specific band was detected for hSTING/TMEM173 antibody (NBP3-18816) at approximately 44 kDa (as indicated) using 10 ug/mL of hSTING/TMEM173 antibody. This experiment was conducted under reducing conditions.




Western blot shows lysates of THP-1 cell line and STING/TMEM173 knockout THP-1 cell line (KO). Nitrocellulose membrane was probed with STING/TMEM173 Antibody (Catalog # NBP3-18816) followed by HRP-conjugated secondary antibody. A specific band was detected for STING/TMEM173 at approximately 41 kDa (as indicated) in the parental THP-1 cell line, but is not detectable in knockout THP-1 cell line. Primary antibody dilution used: 1/1000. The Ponceau stained transfer of the blot is shown. This experiment was conducted under reducing conditions. Image, protocol, and testing courtesy of YCharOS Inc. See ycharos.com for additional details.



PMA-treated THP-1 lysates were prepared and immunoprecipitation was performed using 2.0  $\mu$ g of STING/TMEM173 Antibody (Catalog # NBP3-18816) pre-coupled to Dynabeads protein A. Immunoprecipitated STING/TMEM173 was detected with NBP3-18816. For western blot, NBP3-18816 was used at 1/1000. The Ponceau stained transfer of the blot is shown. SM=4% starting material; UB=4% unbound fraction; IP=immunoprecipitate; HC=antibody heavy chain. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).



PMA-treated THP-1 WT and STING/TMEM173 KO cells were labelled with a green or a far-red fluorescent dye, respectively. Cells were stained with STING/TMEM173 Antibody (Catalog # NBP3-18816) and with an Alexa-fluor 555 coupled secondary antibody including DAPI. Acquisition of the blue (nucleus-DAPI), green (identification of WT cells), red (antibody staining) and far-red (identification of KO cells) channels was performed. Representative images of the blue and red (grayscale) channels are shown. WT and KO cells are outlined with green and magenta dashed line, respectively. Primary antibody dilution used: 1/1000. Image, protocol and testing courtesy of YCharOS Inc. (ycharos.com).





### **Novus Biologicals USA**

10730 E. Briarwood Avenue  
Centennial, CO 80112  
USA  
Phone: 303.730.1950  
Toll Free: 1.888.506.6887  
Fax: 303.730.1966  
nb-customerservice@bio-techne.com

### **Bio-Techne Canada**

21 Canmotor Ave  
Toronto, ON M8Z 4E6  
Canada  
Phone: 905.827.6400  
Toll Free: 855.668.8722  
Fax: 905.827.6402  
canada.inquires@bio-techne.com

### **Bio-Techne Ltd**

19 Barton Lane  
Abingdon Science Park  
Abingdon, OX14 3NB, United Kingdom  
Phone: (44) (0) 1235 529449  
Free Phone: 0800 37 34 15  
Fax: (44) (0) 1235 533420  
info.EMEA@bio-techne.com

### **General Contact Information**

www.novusbio.com  
Technical Support: nb-technical@bio-techne.com  
Orders: nb-customerservice@bio-techne.com  
General: novus@novusbio.com

### **Products Related to NBP3-18816**

---

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
NBP2-24891	Rabbit IgG Isotype Control
NBP2-24683PEP	STING/TMEM173 Antibody Blocking Peptide

---

### **Limitations**

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit [www.novusbio.com/guarantee](http://www.novusbio.com/guarantee)

Earn gift cards/discounts by submitting a review: [www.novusbio.com/reviews/submit/NBP3-18816](http://www.novusbio.com/reviews/submit/NBP3-18816)

Earn gift cards/discounts by submitting a publication using this product:  
[www.novusbio.com/publications](http://www.novusbio.com/publications)

