Product Datasheet

NCAM-1/CD56 Antibody (NCAM1/8392R) NBP3-20529-100ug

Unit Size: 100 ug

Store at 4C. Do not freeze.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-20529

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-20529



NCAM-1/CD56 Antibody (NCAM1	/8392R)	
Product Information		
Unit Size	100 ug	
Concentration	0.2 mg/ml	
Storage	Store at 4C. Do not freeze.	
Clonality	Monoclonal	
Clone	NCAM1/8392R	
Preservative	0.05% Sodium Azide	
Isotype	IgG Kappa	
Purity	Protein A or G purified	
Buffer	10mM PBS with 0.05% BSA	
Product Description		
Description	Positive Controls: Human colon or small intestine. Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.	
Host	Rabbit	
Gene ID	4684	
Oana Cumak al	NICA BAA	


Host	Rabbit
Gene ID	4684
Gene Symbol	NCAM1
Species	Human
Specificity/Sensitivity	This monoclonal antibody reacts with an extracellular domain (close to transmembrane) of CD56/NCAM Anti-CD56 plays an important role in the diagnosis of nodal and pasal NK/T-cell lymphomas

	diagnosis of nodal and hasal Nrv 1-cell lymphomas
_	Recombinant fragment (around aa600-800) of human NCAM-1/CD56 protein (exact sequence is proprietary)

	(construction of properties)
Product Application Details	
Applications	Immunohistochemistry-Paraffin
Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml
Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.

Images

Formalin-fixed, paraffin-embedded human colon stained with NCAM-1/CD56 antibody (NCAM1/8392R). HIER: Tris/EDTA, pH9.0, 45min. Secondary: HRP-polymer, 30min. DAB, 5min.





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112

USA

Phone: 303.730.1950 Toll Free: 1.888.506.6887

Fax: 303.730.1966

nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6

Canada

Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402

canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449

Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com

Technical Support: nb-technical@bio-

techne.com

Orders: nb-customerservice@bio-techne.com

General: novus@novusbio.com

Products Related to NBP3-20529-100ug

HAF008 Goat anti-Rabbit IgG Secondary Antibody [HRP]

NB7160 Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]

NBP3-07122-50ug Recombinant Human NCAM-1/CD56 His Protein

210-TA-005 TNF-alpha [Unconjugated]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-20529

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

