Product Datasheet

COX-2 Antibody (COX2/7803R) - Azide and BSA Free NBP3-21139

Unit Size: 100 ug

Store at -20 to -70C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-21139

Updated 7/16/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-21139



NBP3-21139

COX-2 Antibody (COX2/7803R) - Azide and BSA Free


of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.		
Concentration 1 mg/ml Storage Store at -20 to -70C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone COX2/7803R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Applications Applications Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
Storage Store at -20 to -70C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone COX2/7803R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug
Clonality Monoclonal Clone COX2/7803R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Recommended Dilutions Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	1 mg/ml
CloneCOX2/7803RPreservativeNo PreservativeIsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBSProduct DescriptionPositive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-Paraffin Recommended DilutionsApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Storage	Store at -20 to -70C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
IsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBSProduct DescriptionPositive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-Paraffin Recommended DilutionsApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clone	COX2/7803R
Purity Protein A or G purified Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	No Preservative
Buffer 10mM PBS Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2 ug/ml Application Notes Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG Kappa
Product Description Positive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required. Host Rabbit Gene ID 5743 Gene Symbol PTGS2 Species Human Immunogen Recombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary) Product Application Details Applications Application Notes Immunohistochemistry-Paraffin 1-2 ug/ml Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
DescriptionPositive Controls: Human colon or lung carcinoma tissue (IHC). Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10mM PBS
Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to -80C. Non-hazardous. No MSDS required.HostRabbitGene ID5743Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID5743Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Antibody with azide - store at 2 to 8C. Antibody without azide - store at -20 to
Gene SymbolPTGS2SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Rabbit
SpeciesHumanImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	5743
ImmunogenRecombinant fragment (around aa400-604) of human COX-2 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	PTGS2
sequence is proprietary)Product Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
ApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	\mathbf{v}
Recommended DilutionsImmunohistochemistry-Paraffin 1-2 ug/mlApplication NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application NotesImmunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry-Paraffin
of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2 ug/ml
	Application Notes	Immunohistochemistry (Formalin-fixed): 1-2ug/ml for 30 minutes. at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes. Optimal dilution for a specific application should be determined.



Images

Formalin-fixed, paraffin-embedded human prostate stained with COX-2 antibody (COX2/7803R). HIER: Tris/EDTA, pH9.0, 45min. Secondary: HRP-polymer, 30min. DAB, 5min.

Formalin-fixed, paraffin-embedded human prostate stained with COX-2 antibody (COX2/7803R). HIER: Tris/EDTA, pH9.0, 45min. Secondary: HRP-polymer, 30min. DAB, 5min.







Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-21139

210-TA-005	TNF-alpha [Unconjugated]
NBP1-85499PEP	COX-2 Recombinant Protein Antigen
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-21139

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

