## **Product Datasheet**

### CD63 Antibody (LAMP3/8678R) NBP3-23647-100ug

Unit Size: 100 ug

Store at 4C.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-23647

Updated 7/16/2024 v.20.1

# Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-23647



#### NBP3-23647-100ug

CD63 Antibody (LAMP3/8678R)

Product Information     Unit Size   100 ug     Concentration   0.2 mg/ml     Storage   Store at 4C.     Clonality   Monoclonal     Clone   LAMP3/8678R     Preservative   0.05% Sodium Azide     Isotype   IgG Kappa     Purity   Protein A or G purified     Buffer   10mM PBS with 0.05% BSA     Product Description   Store at 2 to 8C. Non-hazardous. No MSDS required.     Host   Rabbit     Gene ID   967     Gene Symbol   CD63     Species   Human     Marker   Late Endosomes Marker     Species   Human     Marker   Late Endosomes Marker     Specificity/Sensitivity   This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.     Immunogen   Recombinant full-length human CD63 protein     Application Details   Immunohistochemistry-Paraffin     Recombinant full-length further 1-2ug/ml   Application Sint further 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, p	, , , , , , , , , , , , , , , , , , ,	
Concentration0.2 mg/mlStorageStore at 4C.ClonalityMonoclonalCloneLAMP3/8678RPreservative0.05% Sodium AzideIsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBS with 0.05% BSAProduct DescriptionStore at 2 to 8C. Non-hazardous. No MSDS required.HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpeciesSpitopi is epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Information	
Storage Store at 4C.   Clonality Monoclonal   Clone LAMP3/8678R   Preservative 0.05% Sodium Azide   Isotype IgG Kappa   Purity Protein A or G purified   Buffer 10mM PBS with 0.05% BSA   Product Description Store at 2 to 8C. Non-hazardous. No MSDS required.   Host Rabbit   Gene ID 967   Gene Symbol CD63   Species Human   Marker Late Endosomes Marker   Species Human   Marker Late Endosomes Marker   Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.   Immunogen Recombinant full-length human CD63 protein   Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml   Applications Immunohistochemistry-Paraffin 1-2ug/ml   Application Notes Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Unit Size	100 ug
ClonalityMonoclonalCloneLAMP3/8678RPreservative0.05% Sodium AzideIsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBS with 0.05% BSAProduct DescriptionStore at 2 to 8C. Non-hazardous. No MSDS required.HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 22u/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Concentration	0.2 mg/ml
CloneLAMP3/8678RPreservative0.05% Sodium AzideIsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBS with 0.05% BSAProduct DescriptionStore at 2 to 8C. Non-hazardous. No MSDS required.HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-Paraffin 1-2ug/mlApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml	Storage	Store at 4C.
Preservative 0.05% Sodium Azide   Isotype IgG Kappa   Purity Protein A or G purified   Buffer 10mM PBS with 0.05% BSA   Product Description Store at 2 to 8C. Non-hazardous. No MSDS required.   Host Rabbit   Gene ID 967   Gene Symbol CD63   Species Human   Marker Late Endosomes Marker   Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.   Immunogen Recombinant full-length human CD63 protein   Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml   Applications Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG Kappa   Purity Protein A or G purified   Buffer 10mM PBS with 0.05% BSA   Product Description Extra to the state of th	Clone	LAMP3/8678R
Purity Protein A or G purified   Buffer 10mM PBS with 0.05% BSA   Product Description Store at 2 to 8C. Non-hazardous. No MSDS required.   Host Rabbit   Gene ID 967   Gene Symbol CD63   Species Human   Marker Late Endosomes Marker   Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.   Immunogen Recombinant full-length human CD63 protein   Product Application Details Immunohistochemistry-Paraffin   Applications Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Preservative	0.05% Sodium Azide
Buffer10mM PBS with 0.05% BSAProduct DescriptionStore at 2 to 8C. Non-hazardous. No MSDS required.HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG Kappa
Product Description Store at 2 to 8C. Non-hazardous. No MSDS required.   Description Store at 2 to 8C. Non-hazardous. No MSDS required.   Host Rabbit   Gene ID 967   Gene Symbol CD63   Species Human   Marker Late Endosomes Marker   Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.   Immunogen Recombinant full-length human CD63 protein   Product Application Details Immunohistochemistry-Paraffin   Applications Immunohistochemistry-Paraffin 1-2ug/ml   Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
DescriptionStore at 2 to 8C. Non-hazardous. No MSDS required.HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Buffer	10mM PBS with 0.05% BSA
HostRabbitGene ID967Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID 967   Gene Symbol CD63   Species Human   Marker Late Endosomes Marker   Specificity/Sensitivity This monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.   Immunogen Recombinant full-length human CD63 protein   Product Application Details Immunohistochemistry-Paraffin   Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml   Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Store at 2 to 8C. Non-hazardous. No MSDS required.
Gene SymbolCD63SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Rabbit
SpeciesHumanMarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	967
MarkerLate Endosomes MarkerSpecificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	CD63
Specificity/SensitivityThis monoclonal antibody recognizes protein of 26kDa-60kDa, which is identified as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application Details ApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
as CD63. Its epitope is different from that of monoclonal antibody LAMP3/529.ImmunogenRecombinant full-length human CD63 proteinProduct Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Late Endosomes Marker
Product Application Details   Applications Immunohistochemistry-Paraffin   Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml   Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Specificity/Sensitivity	
ApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	Recombinant full-length human CD63 protein
Recommended Dilutions Immunohistochemistry-Paraffin 1-2ug/ml   Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry-Paraffin
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2ug/ml
	Application Notes	formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.

www.novusbio.com





#### Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

#### **Bio-Techne Canada**

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

#### **Bio-Techne Ltd**

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

#### **General Contact Information**

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

#### Products Related to NBP3-23647-100ug

HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
H00000967-G01-2ug	Recombinant Human CD63 Protein
203-IL-010	IL-3 [Unconjugated]

#### Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-23647

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

