Product Datasheet

CD20 Antibody (MS4A1/6993R) [Alexa Fluor® 532] NBP3-24154AF532

Unit Size: 0.1 ml

Store at 4C in the dark.

www.novusbio.com

technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24154AF532

Updated 11/18/2023 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-24154AF532



NBP3-24154AF532

CD20 Antibody (MS4A1/6993R) [Alexa Fluor® 532]

Product Information Unit Size 0.1 ml Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone MS4A1/6993R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 532 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Host Gene ID 931 Gene Symbol MS4A1 Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and P cell blood and Vmphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD2-positive T-cell spherobyre in nown lymphoid tissues. Rarely, CD2-positive T-cell spheromas. In lymphodi tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotyte protein drived lymphorid tissues. Rarely, CD2-positive T-cell spherotyte in known lymphoid tissues. Rarely, CD2-positive T-cell lymphonyte in scases of Hodgkin s disease, partic	J () L	-
Concentration Please see the vial label for concentration. If unlisted please contact technical services. Storage Store at 4C in the dark. Clonality Monoclonal Clone MS4A1/6993R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 532 Purity Protein A purified Buffer SomM Sodium Borate Product Description MS4A1 Host Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Reactivity Notes Expected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee. Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and ymphoid tissues, Rarely, CD20-positive T-cell lymphomas. In lymphoid tissues, Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa 97-297) corresponding to C-terminal of	Product Information	
services. Storage Store at 4C in the dark. Clonality Monoclonal Clone MS4A1/6993R Preservative 0.05% Sodium Azide Isotype IgG Kappa Conjugate Alexa Fluor 532 Purity Protein A purified Buffer 50mM Sodium Borate Product Description Rabbit Gene ID 931 Gene Symbol MS4A1 Species Human Reactivity Notes Expected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee. Specificity/Sensitivity This monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell phenotype in known lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody reascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type. Immunogen Recombinant fragment (around aa 97-297) corresponding to C-terminal of	Unit Size	0.1 ml
ClonalityMonoclonalCloneMS4A1/6993RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B Jymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell localization in the derived lymphomas have been reported. Reactivity has also been noted with Reed-Stemberg cells in cases of Hodgkin s disease, particularly relative. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. <t< th=""><th>Concentration</th><th>•</th></t<>	Concentration	•
CloneMS4A1/6993RPreservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionRabbitHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Storage	Store at 4C in the dark.
Preservative0.05% Sodium AzideIsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Clonality	Monoclonal
IsotypeIgG KappaConjugateAlexa Fluor 532PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionFroduct DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and B cell blood and Jymphoid tissues, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell Iphenotype in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Clone	MS4A1/6993R
ConjugateAlexa Fluor 532PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Preservative	0.05% Sodium Azide
PurityProtein A purifiedBuffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Isotype	IgG Kappa
Buffer50mM Sodium BorateProduct DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell lymphomas. In lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Conjugate	Alexa Fluor 532
Product DescriptionHostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Purity	Protein A purified
HostRabbitGene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Buffer	50mM Sodium Borate
Gene ID931Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Product Description	
Gene SymbolMS4A1SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Host	Rabbit
SpeciesHumanReactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Gene ID	931
Reactivity NotesExpected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Gene Symbol	MS4A1
Specificity/SensitivityThis monoclonal antibody can be used for immunophenotyping of leukemia and malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Species	Human
malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly of lymphocyte predominant type.ImmunogenRecombinant fragment (around aa 97-297) corresponding to C-terminal of	Reactivity Notes	Expected to react with Monkey, Baboon, Gorilla, Dog and Chimpanzee.
	Specificity/Sensitivity	malignant cells, B lymphocyte detection in peripheral blood and B cell localization in tissues. It reacts with the majority of B-cells present in peripheral blood and lymphoid tissues and their derived lymphomas. In lymphoid tissue, germinal center blasts and B-immunoblasts are particularly reactive. It is a reliable antibody for ascribing a B-cell phenotype in known lymphoid tissues. Rarely, CD20-positive T-cell lymphomas have been reported. Reactivity has also been noted with Reed-Sternberg cells in cases of Hodgkin s disease, particularly
	Immunogen	



Notes	Alexa Fluor (R) products are provided under an intellectual property license from Life Technologies Corporation. The purchase of this product conveys to the buyer the non-transferable right to use the purchased product and components of the product only in research conducted by the buyer (whether the buyer is an academic or for-profit entity). The sale of this product is expressly conditioned on the buyer not using the product or its components, or any materials made using the product or its components, in any activity to generate revenue, which may include, but is not limited to use of the product or its components: (i) in manufacturing; (ii) to provide a service, information, or data in return for payment; (iii) for therapeutic, diagnostic or prophylactic purposes; or (iv) for resale, regardless of whether they are resold for use in research. For information on purchasing a license to this product for purposes other than as described above, contact Life Technologies Corporation, 5791 Van Allen Way, Carlsbad, CA 92008 USA or outlicensing@lifetech.com. This conjugate is made on demand. Actual recovery may vary from the stated volume of this product. The volume will be greater than or equal to the unit size stated on the datasheet.
Product Application Details	
Applications	Western Blot, Immunohistochemistry-Paraffin
Recommended Dilutions	Western Blot, Immunohistochemistry-Paraffin
Application Notes	Optimal dilution of this antibody should be experimentally determined.
4	





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-24154AF532

H00000931-P01-2ug	Recombinant Human CD20 GST (N-Term) Protein
210-TA-005	TNF-alpha [Unconjugated]
NBL1-13312	CD20 Overexpression Lysate
NBP1-19371	CD4 Antibody - BSA Free

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24154AF532

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

