Product Datasheet

CD31/PECAM-1 Antibody (C31/8828R) - Azide and BSA Free NBP3-24156

Unit Size: 100 ug

Store at -20 to -80C. Avoid freeze-thaw cycles.

www.novusbio.com



technical@novusbio.com

Protocols, Publications, Related Products, Reviews, Research Tools and Images at: www.novusbio.com/NBP3-24156

Updated 9/11/2024 v.20.1

Earn rewards for product reviews and publications.

Submit a publication at www.novusbio.com/publications Submit a review at www.novusbio.com/reviews/destination/NBP3-24156



NBP3-24156

CD31/PECAM-1 Antibody (C31/8828R) - Azide and BSA Free

Product Information Unit Size 100 ug Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone C31/8828R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and senstive for vascular endothelial cells shinin make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around as 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2u/ml Application Notes I		
Concentration 1 mg/ml Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone C31/8828R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, beingn, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Application Notes Immunohistochemistry-Paraffin 1-2ug/ml<	Product Information	
Storage Store at -20 to -80C. Avoid freeze-thaw cycles. Clonality Monoclonal Clone C31/8828R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Immunohistochemistry-Paraffin Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2ug/ml	Unit Size	100 ug
Clonality Monoclonal Clone C31/8828R Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Daraffin 1-2ug/ml	Concentration	1 mg/ml
CloneC31/8828RPreservativeNo PreservativeIsotypeIgG KappaPurityProtein A or G purifiedBuffer10mM PBSProduct DescriptionAntibody without azide - store at -20 to -80C.HostRabbitGene ID5175Gene SymbolPECAM1SpeciesHumanMarkerEndothelial Cell MarkerSpecificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells which make up blood vescel lining. The level of CD31 expression can help to determine the degree of tumor and gogenesis, and a high level of CD31 expression can belp to determine the degree of tumor and gogenesis, and a high level of CD31 expression can selp to determine the degree of tumor and gogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenImmunohistochemistry-Paraffin (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml	Storage	Store at -20 to -80C. Avoid freeze-thaw cycles.
Preservative No Preservative Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells witch marke up blood vessel lining. The level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clonality	Monoclonal
Isotype IgG Kappa Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel liming. The level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml Applications Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Clone	C31/8828R
Purity Protein A or G purified Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml	Preservative	No Preservative
Buffer 10mM PBS Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Isotype	IgG Kappa
Product Description Antibody without azide - store at -20 to -80C. Host Rabbit Gene ID 5175 Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin 1-2ug/ml Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Purity	Protein A or G purified
DescriptionAntibody without azide - store at -20 to -80C.HostRabbitGene ID5175Gene SymbolPECAM1SpeciesHumanMarkerEndothelial Cell MarkerSpecificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin 1-2ug/ml	Buffer	10mM PBS
HostRabbitGene ID5175Gene SymbolPECAM1SpeciesHumanMarkerEndothelial Cell MarkerSpecificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-Paraffin 1-2ug/mlApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Description	
Gene ID5175Gene SymbolPECAM1SpeciesHumanMarkerEndothelial Cell MarkerSpecificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Description	Antibody without azide - store at -20 to -80C.
Gene Symbol PECAM1 Species Human Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Applications Immunohistochemistry-Paraffin 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Host	Rabbit
SpeciesHumanMarkerEndothelial Cell MarkerSpecificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene ID	5175
Marker Endothelial Cell Marker Specificity/Sensitivity Anti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence. Immunogen Recombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary) Product Application Details Immunohistochemistry-Paraffin Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml Application Notes Immunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Gene Symbol	PECAM1
Specificity/SensitivityAnti-CD31 has shown to be highly specific and sensitive for vascular endothelial cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinApplicationsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Species	Human
cells. Staining of nonvascular tumors (excluding hematopoietic neoplasms) is rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a predictor of tumor recurrence.ImmunogenRecombinant fragment (around aa 538-738) of human CD31/PECAM-1 protein (exact sequence is proprietary)Product Application DetailsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Marker	Endothelial Cell Marker
(exact sequence is proprietary)Product Application DetailsApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Specificity/Sensitivity	rare. CD31 monoclonal antibody reacts with normal, benign, and malignant endothelial cells which make up blood vessel lining. The level of CD31 expression can help to determine the degree of tumor angiogenesis, and a high level of CD31 expression may imply a rapidly growing tumor and potentially a
ApplicationsImmunohistochemistry-ParaffinRecommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Immunogen	
Recommended DilutionsImmunohistochemistry-Paraffin 1-2ug/mlApplication NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Product Application Details	
Application NotesImmunohistochemistry-Paraffin: 1-2ug/ml for 30 minutes at RT. Staining of formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Applications	Immunohistochemistry-Paraffin
formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.	Recommended Dilutions	Immunohistochemistry-Paraffin 1-2ug/ml
	Application Notes	formalin-fixed tissues requires heating tissue sections in 10mM Tris with 1mM EDTA, pH 9.0, for 45 min at 95C followed by cooling at RT for 20 minutes.

www.novusbio.com





Novus Biologicals USA

10730 E. Briarwood Avenue Centennial, CO 80112 USA Phone: 303.730.1950 Toll Free: 1.888.506.6887 Fax: 303.730.1966 nb-customerservice@bio-techne.com

Bio-Techne Canada

21 Canmotor Ave Toronto, ON M8Z 4E6 Canada Phone: 905.827.6400 Toll Free: 855.668.8722 Fax: 905.827.6402 canada.inquires@bio-techne.com

Bio-Techne Ltd

19 Barton Lane Abingdon Science Park Abingdon, OX14 3NB, United Kingdom Phone: (44) (0) 1235 529449 Free Phone: 0800 37 34 15 Fax: (44) (0) 1235 533420 info.EMEA@bio-techne.com

General Contact Information

www.novusbio.com Technical Support: nb-technical@biotechne.com Orders: nb-customerservice@bio-techne.com General: novus@novusbio.com

Products Related to NBP3-24156

210-TA-005	TNF-alpha [Unconjugated]
NBP2-54655PEP	CD31/PECAM-1 Recombinant Protein Antigen
NB7160	Goat anti-Rabbit IgG (H+L) Secondary Antibody [HRP]
HAF008	Goat anti-Rabbit IgG Secondary Antibody [HRP]

Limitations

This product is for research use only and is not approved for use in humans or in clinical diagnosis. Primary Antibodies are guaranteed for 1 year from date of receipt.

For more information on our 100% guarantee, please visit www.novusbio.com/guarantee

Earn gift cards/discounts by submitting a review: www.novusbio.com/reviews/submit/NBP3-24156

Earn gift cards/discounts by submitting a publication using this product: www.novusbio.com/publications

www.novusbio.com

